

Computation of the MinMax Measure

S. Munirathnam H.K. Kesavan

Peter Roe

Department of Systems Design Engineering,
University of Waterloo,
Waterloo, Ontario, N2L 3G1, Canada.

August, 1998

Abstract

The MinMax measure of information, defined by Kapur, Baciú and Kesavan [6], is a quantitative measure of the information contained in a given set of moment constraints. It is based on both maximum and minimum entropy. Computational difficulties in the determination of minimum entropy probability distributions (MinEPD) have inhibited exploration of the full potential of minimum entropy and, hence, the MinMax measure. Initial attempts to solve the minimum entropy problem were directed towards finding analytical solutions for some specific set of constraints. Here, we present a numerical solution to the general minimum entropy problem and discuss the significance of minimum entropy and the MinMax measure. Some numerical examples are given for illustration.

1 Introduction

In this paper the problem of estimating the amount of information contained in a given set of moment constraints is considered.

Let X be a discrete random-variate of a probabilistic system which takes values from $\{x_1, x_2, \dots, x_n\}$ and the corresponding probability distribution be $\mathbf{p} = (p_1, p_2, \dots, p_n)$. Suppose the only information available about the random-variate X is that it satisfies the natural constraint of probabilities

$$\sum_{i=1}^n p_i = 1, \quad p_i \geq 0, \quad i = 1, 2, \dots, n \quad (1)$$

and the moment constraints

$$\sum_{i=1}^n g_{ri} p_i = a_r, \quad r = 1, 2, \dots, m, \quad (2)$$

where g_{ri} and a_r are known constants. Our goal is to measure the amount of information provided by the constraints (1) and (2) about the probabilistic system.

One possible method to measure the amount of information contained in a given set of constraints is suggested by Jaynes' maximum entropy principle (MaxEnt). We first maximize the Shannon entropy measure

$$S(\mathbf{p}) = - \sum_{i=1}^n p_i \ln p_i \quad (3)$$

subject to (1) only, to get the uniform distribution, $U = (\frac{1}{n}, \frac{1}{n}, \dots, \frac{1}{n})$, as the maximum entropy probability distribution (MaxEPD) with entropy value $\ln n$. The maximum entropy probability distribution is the most unbiased, most random and most uniform probability distribution in the light of the information available. We next maximize (3) subject to (1) and (2) to obtain another MaxEPD with entropy value S_{max} .

In the presence of additional information, it is known that uncertainty decreases and is less than what is contained in the uniform distribution. Hence the Shannon entropy, being a measure of uncertainty in a probability distribution, decreases (at least does not increase) and as such $S_{max} \leq \ln n$. The difference in the maximum entropy values is due to the additional constraints and hence, it is suggested that

$$I_1 \stackrel{\text{def}}{=} \ln n - S_{max} \quad (4)$$

can be used as a measure of information provided by the constraints (2). This has been used in some problems as a measure of information contained in moment constraints [3, 14].

The drawback in the definition given in equation (4) is that I_1 represents a reduction in *maximum uncertainty* and not a reduction in *uncertainty*, due to the presence of additional constraints. This difference is illustrated using an example of the dice problem [6]. Given some information about the throws of a die, the problem is to assign a probability distribution consistent with the available information. Suppose we have no information about a six-faced die except that provided by (1). The MaxEPD is the uniform distribution and the maximum entropy value is $\ln 6$. Now, if the average score of throws of the die is given to be 3.5, the MaxEPD is still the uniform distribution and the maximum entropy value is $\ln 6$. This implies $I_1 = 0$, that is, the mean value provides no information about the characteristics of the die. Since I_1 measures reduction in maximum uncertainty, it provides only a lower bound for the information contained in the constraints.

When the available information is only the natural constraint, all probability distributions satisfy the given constraint. In the presence of more information in the form of moment constraints, the choice of probability distributions is restricted. Thus, any measure of information provided by a set of constraints should be a measure of the degree of restriction placed on the choice of probability distribution. Kapur, Baciú, and Kesavan [6] used the concept of minimum entropy to define such a measure of information.

2 Minimum Entropy and the MinMax Measure

If we are given only the natural constraint (1), the MaxEPD is the uniform distribution with entropy value $\ln n$. The minimum entropy probability distribution (MinEPD) is one




Figure 1: Variation of S_{max} and S_{min} with constraints

of the n degenerate distributions $(1, 0, \dots, 0)$, $(0, 1, \dots, 0)$, \dots , $(0, 0, \dots, 1)$ with entropy value 0. With only the natural constraint, the minimum entropy distributions are the n distinct possible outcomes corresponding to the n distinct values of the random-variate X . The MinEPD gives the most biased, the least random and the least uniform probability distribution subject to the available information. The minimum entropy has the potential to “... provide the key to the understanding of the information theoretic structure(s) which lie dormant in a probabilistic system” [9]. For this reason, Watanabe [16] described pattern recognition as a quest for minimum entropy.

With additional moment constraints, the choice of probability distributions is reduced. A restricted set of probability distributions has a smaller maximum entropy than the original set and it has also, in general, a larger minimum entropy value. Thus each additional piece of information in the form of constraints on p_i results in a decrease (or no change) in S_{max} and an increase (or no change) in S_{min} . As indicated in Fig. 1, S_{max} is a monotonic decreasing function and S_{min} is a monotonic increasing function.

In general, every consistent additional constraint reduces

$$I = S_{max} - S_{min}. \quad (5)$$

This quantity is called the *uncertainty gap*. The greater the information contained in a constraint, the greater will be the reduction in S_{max} , the greater will be the increase in S_{min} , and the greater is the reduction in the uncertainty gap.

*The reduction in uncertainty gap due to a constraint set is called the **MinMax measure of information** contained in the constraint set.*

The MinMax measure is a relative measure. For two constraint sets C_1 and C_2 , the MinMax measure of C_2 with respect to C_1 is given in (6) as the difference of uncertainty gap

of C_2 from that of C_1 .

$$I(C_1; C_2) = [S_{max}(C_1) - S_{min}(C_1)] - [S_{max}(C_2) - S_{min}(C_2)] \quad (6)$$

For any constraint set C_1 , the MinMax measure is evaluated with respect the natural constraints and is given by

$$I(C_1) = [S_{max}(N) - S_{min}(N)] - [S_{max}(C_1) - S_{min}(C_1)]. \quad (7)$$

In the case of discrete random-variate X the MinMax measure for a constraint set C is given by

$$I(C) = \ln n - [S_{max}(C) - S_{min}(C)]. \quad (8)$$

3 An Algorithm for the MinMax measure

The calculation of the MinMax measure for a given set of constraints, C , involves calculating the maximum and minimum entropy values conditioned upon the given moment constraints. Although our method generates the maximum and minimum entropy probability distributions to calculate the entropy values, there should be other methods to obtain the entropy values directly.

The computation of the maximum entropy and the MaxEPD is a concave maximization problem. Analytical solutions can be obtained using the Lagrange multiplier method [2, 8]. Accordingly, the maximization of the Shannon entropy measure subject to the constraints (1) and (2) gives

$$p_i = \exp(-\lambda_0 - \lambda_1 g_{1i} - \lambda_2 g_{2i} - \dots - \lambda_m g_{mi}); \quad i = 1, 2, \dots, n. \quad (9)$$

where $\lambda_0, \lambda_1, \dots, \lambda_m$ are Lagrange multipliers corresponding to the $m + 1$ constraints. The Lagrange multipliers are determined by substituting for p_i from (9) in (1) and (2) and simplifying to get

$$\lambda_0 = \ln\left(\sum_{i=1}^n \exp\left(-\sum_{j=1}^m \lambda_j g_{ji}\right)\right) \quad (10)$$

$$a_r = \frac{\sum_{i=1}^n g_{ri} \exp\left(-\sum_{j=1}^m \lambda_j g_{ji}\right)}{\sum_{i=1}^n \exp\left(-\sum_{j=1}^m \lambda_j g_{ji}\right)}, \quad r = 1, 2, \dots, m \quad (11)$$

Equations (11) form a system of m nonlinear equations in m unknowns. These equations can be represented in a form more convenient for numerical solution as a residual

$$R_r = 1 - \frac{\sum_{i=1}^n g_{ri} \exp\left(-\sum_{j=1}^m \lambda_j g_{ji}\right)}{a_r \sum_{i=1}^n \exp\left(-\sum_{j=1}^m \lambda_j g_{ji}\right)} \quad (12)$$

for $r = 1, 2, \dots, m$.

A solution for the Lagrange multipliers can be obtained using nonlinear programming, by minimizing the sum of the squares of the residuals

$$R = \sum_{r=1}^m R_r^2. \quad (13)$$

In the examples illustrated in section 4, the Levenberg-Marquardt method [2, 15] is used for solving the least squares problem.

Evaluating the minimum entropy value for the given set of constraints is the next step in computing the MinMax measure. The minimum entropy problem is given by

$$\text{global } \min_{p \in D \cap C} S(p) = \sum_{i=1}^n s_i(p_i) = - \sum_{i=1}^n p_i \ln p_i \quad (14)$$

where $C = \prod_{i=1}^n C_i$ with $C_i = [l_i, u_i]$, $l_i, u_i \in [0, 1]$ and $D = \{p : \sum_{i=1}^n p_i = 1, \sum_{i=1}^n g_{ri} p_i = a_r, r = 1, 2, \dots, m\}$.

The minimum entropy problem is a global optimization problem, specifically, a concave minimization problem of a separable objective function. This problem is known to be NP-Hard. Analytical solutions for the minimum entropy problem have been obtained for some specific cases. Kapur, Baciuc and Kesavan [5] have obtained the expression for minimum entropy if the available information is variance, V , of the random variate X which takes values from $\{x_1, x_2, \dots, x_n\}$. Let $x_1 \leq x_2 \leq \dots \leq x_n$. It has been shown in [5] that $0 \leq V \leq \frac{(x_n - x_1)^2}{4}$ and the minimum entropy probability distribution is the one assigning all probability mass on x_1 and x_n . The minimum entropy value is given by

$$S \left(\frac{1}{2} + \sqrt{\frac{1}{4} - \frac{V}{(x_n - x_1)^2}}, \frac{1}{2} - \sqrt{\frac{1}{4} - \frac{V}{(x_n - x_1)^2}} \right) \quad (15)$$

Furthermore, assuming $x_1 \leq x_2 \leq \dots \leq x_n$, for a mean constraint, it is shown [10] that the minimum entropy probability distribution subject to

$$p_1 x_1 + p_2 x_2 + \dots + p_n x_n = \mu, \quad x_1 \leq \mu \leq x_n \quad (16)$$

is concentrated on

1. a single value x_i if $\mu = x_i$,
2. either (x_1, x_i) or (x_{i-1}, x_n) if $x_{i-1} < \mu < x_i$.

The corresponding minimum entropy values are

$$S_{min}(\mu) = \begin{cases} 0 & \text{if } \mu = x_i \\ \min \left\{ S\left(\frac{x_n - \mu}{x_n - x_{i-1}}, \frac{\mu - x_n}{x_n - x_{i-1}}\right), S\left(\frac{x_i - \mu}{x_i - x_1}, \frac{\mu - x_1}{x_i - x_1}\right) \right\} & \text{if } x_{i-1} < \mu < x_i \end{cases} \quad (17)$$

These solutions also indicate the difficulty in extending such analysis to general problems.

Concave minimization problems are important in global optimization due to the structure of their objective functions. Many numerical solution methods are available for concave minimization [1, 4]. An important property of concave minimization is that the global minimum point is always found at an extremum point of the feasible set. For this reason, linear programming is an essential part of any computational algorithm to solve this problem.

Here we present a numerical solution to the general minimum entropy problem which is based on an algorithm suggested by Phillips and Rosen [13] for constrained minimization of separable concave functions. They use a rectangular successive partitioning approach for solving this concave minimization problem [1]. Indirect information, in the form of lower bounds to the objective function values, is used to decide whether or not the global minimum is contained in the partition of interest. Additionally, some sufficient conditions are used for recognizing a local minimum as a global optimal solution for the problem. The details of the algorithm and its implementation are discussed in [13, 12]. Some salient steps of the algorithm are given below.

Initialization

The problem constraints are preprocessed to obtain an initial hyperrectangle C^0 for the feasible set of solutions. This can be obtained by defining a feasible interval $[l_i^0, u_i^0]$ for each univariate p_i . The minimum l_i^0 is obtained by solving the linear program $\min p_i$ subject to $p \in D \cap [0, 1]^n$ for $i = 1, 2, \dots, n$. Similarly, the maximum u_i^0 can be obtained by solving the linear program $\max p_i$ subject to $p \in D \cap [0, 1]^n$ for $i = 1, 2, \dots, n$.

The set of subproblems, S , is initialized to $\min S(p)$ subject to $p \in D \cap C^0$. Let $k \leftarrow 0$. At each iteration k of the algorithm the following steps are performed.

Bounding

Select a subproblem, s_k , defined as $\min S(p)$ subject to $p \in D \cap C^{s_k}$ from the list S of currently open subproblems. The bounds for each concave subproblem s_k is determined by solving a linear programming relaxation. For each univariate concave term $s_i(p_i)$ in the objective, a linear underestimating function, $\gamma_i(p_i)$, is constructed such that it intersects $s_i(p_i)$ at the current bounds $l_i^{s_k}$ and $u_i^{s_k}$. It is well known that $\gamma_i(p_i)$ is a convex envelope of $s_i(p_i)$ over $[l_i^{s_k}, u_i^{s_k}]$ and that the convex envelope of the separable function $S(p)$ over the rectangular set C^{s_k} is the sum of convex envelope of its individual terms $s_i(p_i)$ over their respective intervals in C^{s_k} . Hence, $\Gamma(p) = \sum_{i=1}^n \gamma_i(p_i)$ is a linear underestimator of $S(p)$ over C^{s_k} .

In the bounding step, the relaxed linear program given by

$$\Gamma(p^{s_k}) = \min_{p \in D \cap C^{s_k}} \Gamma(p) \quad (18)$$

is solved to obtain the minimum p^{s_k} . If p^* is a global optimum solution of the minimum entropy problem, then $\Gamma(p^{s_k}) \leq S^* = S(p^*) \leq S(p^{s_k})$. If $S(p^{s_k}) - \Gamma(p^{s_k}) \leq \epsilon$, a small pre-specified non-negative constant, then p^{s_k} is accepted as a global optimal solution.

If the difference $(S(p^{s_k}) - \Gamma(p^{s_k}))$ exceeds ϵ , the underestimating function $\Gamma(p)$ is refined to obtain a better approximation for $S(p)$ and a solution for the minimum entropy problem.

Kesavan and Zhao [11] use p^{s_k} as a good approximation of the global minimum for MinEPD corresponding to mean and variance constraints.

Refinement

Using the point p^{s_k} a better underestimating function for $S(p)$ over $D \cap C^{s_k}$ is obtained. For each concave univariate p_i , let $\gamma_i^{(1)}(p_i)$ be the linear underestimating function of $s_i(p_i)$ over the interval $[l_i^{s_k}, p_i^{s_k}]$ and let $\gamma_i^{(2)}(p_i)$ be the linear underestimating function of $s_i(p_i)$ over the interval $[p_i^{s_k}, u_i^{s_k}]$.

Since the minimum value occurs at extreme points of the solution set, the solution p^{s_k} minimizing $\Gamma(p)$ has for some i , $p_i = l_i^{s_k}$, or $p_i = u_i^{s_k}$ and others $p_i \in (l_i^{s_k}, u_i^{s_k})$. Assume that $p_i^{s_k} \in (l_i^{s_k}, u_i^{s_k})$ for $i = 1, 2, \dots, n_0$ and $p_i \in \{l_i^{s_k}, u_i^{s_k}\}$ for $i = n_0 + 1, \dots, n$. It is proved [13, Theorem 1] that if p^{s_k} solves the multiple cost row linear program given by

$$\min_{p \in D \cap C^{s_k}} \sum_{i=1}^{n_0} \gamma_i^{j_i}(p_i) + \sum_{n_0+1}^n \gamma_i(p_i) \quad (19)$$

where $j_i \in \{1, 2\}$, for $i = 1, 2, \dots, n_0$, for all 2^{n_0} distinct cost rows and $p^* \in D \cap C^{s_k}$, then p^{s_k} satisfies $S^* = S(p^{s_k})$; that is, p^{s_k} is a global optimal vertex for the concave minimization problem. This implies that if all combinations of underestimating functions have their minimum at p^{s_k} , then p^{s_k} is a global optimal solution.

The check for optimality of p^{s_k} involves 2^{n_0} combinations of cost rows and requires exponential time. Phillips and Rosen [13] identified that an alternate check for optimality can be performed in polynomial time. It uses the fact that a basis remains optimal when the cost coefficients are perturbed, provided that the reduced costs remain non-negative. If this condition is satisfied the p^{s_k} is a global optimal solution. If some of the non-basic variables are sensitive to the perturbation in the cost coefficients, they are used to obtain a better solution than p^{s_k} for the minimization problem.

Let B be the ordered set of indices of the basic variables at the point p^{s_k} . Let N be the set of indices of non-basic variables which do not satisfy the above condition on the cost coefficients. For $i \in N \cup B$, let the points $p^{(1,i)}$ and $p^{(2,i)}$ denote the solution to the pairs of linear programs

$$\min_{p \in D \cap C^{s_k}} \sum_{\substack{j=1 \\ j \neq i}}^n \gamma_j(p_j) + \gamma_i^{(1)}(p_i), \text{ and} \quad (20)$$

$$\min_{p \in D \cap C^{s_k}} \sum_{\substack{j=1 \\ j \neq i}}^n \gamma_j(p_j) + \gamma_i^{(2)}(p_i) \quad (21)$$

respectively. Let

$$\begin{aligned}\Gamma^{(1,i)} &= \sum_{\substack{j=1 \\ j \neq i}}^n \gamma_j(p_j^{(1,i)}) + \gamma_i^{(1)}(p_i^{(1,i)}) \\ \Gamma^{(2,i)} &= \sum_{\substack{j=1 \\ j \neq i}}^n \gamma_j(p_j^{(2,i)}) + \gamma_i^{(2)}(p_i^{(2,i)}) \\ S^{(1,i)} &= S(p^{(1,i)}) \quad \text{and} \quad S^{(2,i)} = S(p^{(2,i)}).\end{aligned}$$

$\Gamma^{(1,i)}$ and $\Gamma^{(2,i)}$ are the minimum values of $\Gamma(p)$ over smaller hyperrectangles obtained by partitioning $D \cap C^{s_k}$ only in the i th dimension. The minimum of $\Gamma^{(1,i)}$ and $\Gamma^{(2,i)}$ is a lower bound of $S(p)$ with respect to the partitioning in the i th dimension. This leads to $\Gamma(p^{s_k}) \leq \Gamma \leq S^* \leq \min\{S, S(p^{s_k})\}$, where $\Gamma = \max_{i \in N \cup B} \{\min\{\Gamma^{(1,i)}, \Gamma^{(2,i)}\}\}$ and $S = \min_{i \in N \cup B} \{S^{(1,i)}, S^{(2,i)}\}$ [13, Theorem 3]. if $S - \Gamma \leq \epsilon$, then $p^{s_k} = \arg \min(S)$ gives a global optimal solution.

This step involves solving $2|N \cup B|$ linear programs. However, it is observed that if the global minimum p^{s_k} is updated by one of these linear programs, it usually corresponds to the one which is most sensitive to the perturbations in the cost coefficient. This heuristic is used in the implementation of the algorithm to obtain a refinement of p^{s_k} .

Branching

If the solution is not obtained by the refinement process, the solution provided by $2|N \cup B|$ linear programs is used to eliminate portions of $D \cap C^{s_k}$ that cannot contain the global optimal solution, p^* . In the branching stage, the subproblem s_k is replaced by a different subproblem with smaller solution space or by replacing the subproblem with two new ones.

Let $i \in N \cup B$, if $\Gamma^{(1,i)} \geq S^{(1,i)}$ the region $C_i^{(1)} = [l_i^{s_k}, p_i^{s_k}] \prod_{j \neq i} [l_j^{s_k}, u_j^{s_k}]$ can be discarded since it can be shown that $p^* \notin C_i^{(1)}$. Similarly, if $\Gamma^{(2,i)} \geq S^{(2,i)}$ the region $p^* \notin C_i^{(2)} = [p_i^{s_k}, u_i^{s_k}] \prod_{j \neq i} [l_j^{s_k}, u_j^{s_k}]$ and hence can be discarded. If both $\Gamma^{(1,i)} \geq S^{(1,i)}$ and $\Gamma^{(2,i)} \geq S^{(2,i)}$, then p^{s_k} is, in fact, the global minimal point. If some subregions are eliminated, the solution space $D \cap C^{s_k}$ is reduced and the iteration is restarted.

If no subregion is eliminated, the solution space $D \cap C^{s_k}$ is partitioned along the edge that corresponds to a variable most responsible for the gap $S(p) - \Gamma(p)$. if $j' = \arg \max_{i=1,2,\dots,n} [s_i(p_i^{s_k}) - \gamma_i(p_i^{s_k})]$, then C^{s_k} is partitioned along l resulting in two new subproblems:

$$\begin{aligned}\min S(p) \text{ subject to } p \in D \cap C^{s_{k1}}, C^{s_{k1}} &= [l_{j'}^{s_k}, p_{j'}^{s_k}] \prod_{j \neq j'} [l_j^{s_k}, u_j^{s_k}] \text{ and} \\ \min S(p) \text{ subject to } p \in D \cap C^{s_{k2}}, C^{s_{k2}} &= [p_{j'}^{s_k}, u_{j'}^{s_k}] \prod_{j \neq j'} [l_j^{s_k}, u_j^{s_k}].\end{aligned}$$

These two problems are added to the set S of subproblems and the iteration continues by incrementing k and selecting a different subproblem.

This method provides an ϵ -approximate solution to the minimum entropy problem. This approximate solution is used to compute the MinMax measure for a constraint set. First, the maximum and minimum entropy values corresponding to the constraint set are computed. The MinMax measure is obtained using the equation (8).

4 Numerical Example: A traffic engineering problem

A problem in traffic engineering is considered here to illustrate the above algorithms for computing the MinMax measure [7]. The computational results in this section were obtained on a SUN Sparc 2 machine. A stopping tolerance of $\epsilon = 0.001$ was used.

Assume that in a city there are six residential areas (or origins) and one business district (or destination). A transportation system should efficiently move people from the origins to the destination. The usual information available about this system is the cost of travel from one origin to the destination, and the average cost of travel to a destination. Let the travel cost from each of these origins to the destination be 1.25, 2.25, 4.25, 5.00, 5.40, 5.75 dollars, respectively. Let the average cost of travel be \$3.5. Based on this information, a transportation planning engineer would like to estimate the percentage of population residing at each of the origins. Also of interest is the amount of knowledge gained about the system from the available information.

Let X be the random-variate for the transportation system which takes the values from $\{1.25, 2.25, 4.25, 5.00, 5.40, 5.75\}$ and the corresponding probability distribution be (p_1, p_2, \dots, p_6) . The given information can be formulated as the following constraint set

$$C = \left\{ \sum_{i=1}^6 p_i = 1; p_i \geq 0, i = 1, 2, \dots, 6; \sum_{i=1}^6 p_i x_i = 3.5 \right\}. \quad (22)$$

With only the natural constraint, the MaxEPD corresponds to the uniform distribution $(\frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6})$ with entropy value 1.791759. The MinEPD is one of the certain distributions $(1, 0, 0, 0, 0, 0)$, $(0, 1, 0, 0, 0, 0)$, $(0, 0, 1, 0, 0, 0)$, $(0, 0, 0, 1, 0, 0)$, $(0, 0, 0, 0, 1, 0)$, $(0, 0, 0, 0, 0, 1)$ with entropy value 0. The MaxEPD with the additional constraint of arithmetic mean is

$$p_{max} = (0.250756 \ 0.212893 \ 0.153455 \ 0.135726 \ 0.127124 \ 0.120045) \quad (23)$$

with entropy value 1.751579. The corresponding MinEPD is obtained as

$$p_{min} = (0.250000 \ 0.000000 \ 0.750000 \ 0.000000 \ 0.000000 \ 0.000000) \quad (24)$$

with entropy value 0.562335. The MinMax measure for the constraint set, C , is

$$I(C) = (1.791759 - 0) - (1.751579 - 0.562335) = 0.602515. \quad (25)$$

One can consider other constraint sets for this problem to identify a set of constraints which contains more information about the probabilistic system. In the following, we analyze the effect of geometric mean constraint on this problem. Let the geometric mean of the cost of travel be g . The given set of constraints is

$$C = \left\{ \sum_{i=1}^6 p_i = 1; p_i \geq 0, i = 1, 2, \dots, 6; \sum_{i=1}^6 p_i \ln(x_i) = \ln g \right\}. \quad (26)$$

The geometric mean $(\ln g)$ takes values from $[\ln 1.25, \ln 5.75]$. The MinEPD for various values of the geometric mean is given in Table 1. The graphs of S_{max} , S_{min} , and the MinMax

measure are given in Fig. 2.

The S_{max} curve is concave with respect to differing geometric mean values. As the x_i values increases from 1.25, the maximum entropy distribution becomes more uniform. Further increasing the geometric mean value pushes the maximum entropy distribution from the uniform distribution and the maximum entropy value decreases. The S_{min} value is piecewise concave with respect to the geometric mean value. The geometric mean values where S_{min} attains local minimum correspond to the certain or degenerate distributions of the random variable. In this case, they are the certain distributions $(1, 0, 0, 0, 0, 0)$, $(0, 1, 0, 0, 0, 0)$, $(0, 0, 1, 0, 0, 0)$, $(0, 0, 0, 1, 0, 0)$, $(0, 0, 0, 0, 1, 0)$, $(0, 0, 0, 0, 0, 1)$ representing the six different x_i values of the random variate. This corresponds to the distinct patterns in the problem.

5 Concluding Remarks

The main contribution of this paper is the numerical method to solve the minimum entropy problem for a general set of constraints to enable the computation of the MinMax measure. Earlier attempts to solve this NP-Hard problem were limited to finding analytical solutions for some simple moments, like the arithmetic mean and variance. Here, a method to obtain an ϵ -approximate solution to the minimum entropy problem using rectangular successive partitioning approach for concave minimization was presented.

The main objective of solving the minimum entropy problem, apart from exploring the properties of minimum entropy, is to compute the MinMax measure for a given set of constraints. The MinMax measure enables us to determine how much information is contained in a set of moment constraints and to discriminate between constraint sets.

Given a number of constraint sets, the MinMax measure will be able to identify the most informative set of moment constraints. This also provides a method for solving inverse problems [12] where the goal is to arrive at a set of moment constraints corresponding to an empirical probability distribution. Of the constraint sets considered, the one with the *maximum* MinMax measure has the most information about the probabilistic system. This set of moment constraints can be identified to “best” characterize the behavior of the system.

The numerical method presented here for the minimum entropy problem uses only the properties of a concave function. The properties of Shannon entropy measure can be used to improve the algorithm. Other global optimization techniques, like stochastic optimization and interval methods, need to be explored for this problem. Faster and accurate solutions to the minimum entropy problem will provide better methods for computing the MinMax measure and enable on-line applications of the MinMax measure.

References

- [1] H.P. Benson. Concave minimization : Theory, applications and algorithms. In Horst. R. and Pardalos. P.M., editors, *Handbook of Global Optimization*. Kluwer Academic Publications, 1995.
- [2] R. Fletcher. *Practical Methods of Optimization*. John Wiley, 1991.

- [3] R.M. Gray and J.E. Shore. Minimum cross-entropy pattern classification and cluster analysis. *IEEE trans. on PAMI*, 4(1):11–17, 1982.
- [4] R. Horst and H. Tuy. *Global Optimization: Deterministic Approaches*. Springer-Verlag, Berlin, 1993.
- [5] J.N. Kapur, G. Baciú, and H.K. Kesavan. On the relationship between variance and minimum entropy. *Internal publication of Univ. of Waterloo, Waterloo, Canada*, 1994.
- [6] J.N. Kapur, G. Baciú, and H.K. Kesavan. The minmax information measure. *Int. J. Systems Sci.*, 26(1):1–12, 1995.
- [7] J.N. Kapur and H.K. Kesavan. *Generalized Maximum Entropy Principle (with Applications)*. Sandford Educational Press, Waterloo, 1989.
- [8] J.N. Kapur and H.K. Kesavan. *Entropy Optimization Principles with Applications*. Academic Press, Inc., New York, 1992.
- [9] H.K. Kesavan and G. Baciú. The role of entropy optimization principles in the study of probabilistic systems. *Proc. Int. Cong. Cybernetics and Systems (ICCS 93)*, pages 25–35, 1993.
- [10] H.K. Kesavan and L. Yuan. Minimum entropy and information measure. *IEEE trans. on SMC*, Aug, 1998.
- [11] H.K. Kesavan and Q. Zhao. An heuristic approach for finding minimum entropy probability distributions. *Internal publication of Univ. of Waterloo, Waterloo, Canada*, 1995.
- [12] S. Munirathnam. The role of minmax entropy measure in probabilistic systems design. Master’s thesis, University of Waterloo, Waterloo, Ontario, Canada, 1998.
- [13] A.T. Phillips and J.B. Rosen. Sufficient conditions for solving linearly constrained separable concave global minimization problems. *J. of Global Optimization*, 3:79–94, 1993.
- [14] S.D. Pietra, V.D. Pietra, and J. Lafferty. Inducing features of random fields. *IEEE trans. on PAMI*, 19(4):380–393, 1997.
- [15] L.E. Scales. *Introduction to Non-linear Optimization*. Macmillan, London, 1985.
- [16] S. Watanabe. Pattern recognition as a quest for minimum entropy. *Pattern Recognition*, 13:381–387, 1981.

$\ln g$	p_1	p_2	p_3	p_4	p_5	p_6	S_{min}
0.223144	1.000000	0.000000	0.000000	0.000000	0.000000	0.000000	0.000000
0.273144	0.967236	0.000000	0.000000	0.000000	0.000000	0.032764	0.144223
0.323144	0.934471	0.000000	0.000000	0.000000	0.000000	0.065529	0.241917
0.373144	0.901707	0.000000	0.000000	0.000000	0.000000	0.098293	0.321316
0.423144	0.868943	0.000000	0.000000	0.000000	0.000000	0.131057	0.388391
0.473144	0.836179	0.000000	0.000000	0.000000	0.000000	0.163821	0.445952
0.523144	0.803415	0.000000	0.000000	0.000000	0.000000	0.196585	0.495632
0.573144	0.770650	0.000000	0.000000	0.000000	0.000000	0.229350	0.538490
0.623144	0.737886	0.000000	0.000000	0.000000	0.000000	0.262114	1.130702
0.673144	0.234415	0.765585	0.000000	0.000000	0.000000	0.000000	0.544556
0.723144	0.149350	0.850650	0.000000	0.000000	0.000000	0.000000	0.421580
0.773144	0.064286	0.935714	0.000000	0.000000	0.000000	0.000000	0.238601
0.823144	0.000000	0.986983	0.000000	0.000000	0.000000	0.013017	0.069445
0.873144	0.000000	0.933693	0.000000	0.000000	0.000000	0.066307	0.243980
0.923144	0.000000	0.880403	0.000000	0.000000	0.000000	0.119597	0.366121
0.973144	0.000000	0.827114	0.000000	0.000000	0.000000	0.172886	0.460433
1.023144	0.000000	0.773824	0.000000	0.000000	0.000000	0.226176	0.534614
1.073144	0.000000	0.720535	0.000000	0.000000	0.000000	0.279465	0.592447
1.123144	0.264571	0.000000	0.735429	0.000000	0.000000	0.000000	0.577784
1.173144	0.223713	0.000000	0.776287	0.000000	0.000000	0.000000	0.531567
1.223144	0.182856	0.000000	0.817144	0.000000	0.000000	0.000000	0.475697
1.250000	0.160911	0.000000	0.839089	0.000000	0.000000	0.000000	0.441177
1.273144	0.141999	0.000000	0.858001	0.000000	0.000000	0.000000	0.408576
1.323144	0.101142	0.000000	0.898858	0.000000	0.000000	0.000000	0.327585
1.373144	0.060285	0.000000	0.939715	0.000000	0.000000	0.000000	0.227751
1.423144	0.019428	0.000000	0.980572	0.000000	0.000000	0.000000	0.095805
1.473144	0.000000	0.000000	0.913243	0.000000	0.000000	0.086757	0.294970
1.523144	0.062248	0.000000	0.000000	0.937752	0.000000	0.000000	0.233109
1.573144	0.026181	0.000000	0.000000	0.973819	0.000000	0.000000	0.121205
1.623144	0.043229	0.000000	0.000000	0.000000	0.956771	0.000000	0.178074
1.673144	0.009059	0.000000	0.000000	0.000000	0.990941	0.000000	0.051631
1.723144	0.017074	0.000000	0.000000	0.000000	0.000000	0.982926	0.086422

Table 1: MinEPD and S_{min} values for geometric mean constraint




Figure 2: S_{max} , S_{min} and MinMax plot for geometric mean constraint