

Markov Network Independencies


Sargur Srihari
srihari@cedar.buffalo.edu

Topics

- Markov Network Independencies
 - Basic Independencies
 - Independencies Revisited
 - From Distributions to Graphs

Markov network captures independencies of interactions

Misconception MN with factors



$$P(a,b,c,d) = \frac{1}{Z} \phi_1(a,b) \cdot \phi_2(b,c) \cdot \phi_3(c,d) \cdot \phi_4(d,a)$$

where

$$Z = \sum_{a,b,c,d} \phi_1(a,b) \cdot \phi_2(b,c) \cdot \phi_3(c,d) \cdot \phi_4(d,a)$$

Factorization of Distribution and Independences

P supports $(X \perp Y|Z)$ iff we can write distribution as $P(X) = \Phi_1(X,Z) \Phi_2(Y,Z)$

$$P(A,B,C,D) = \left[\frac{1}{Z} \phi_1(A,B) \cdot \phi_2(B,C) \right] \phi_3(C,D) \cdot \phi_4(D,A)$$

Factor with $\{B, \{A, C\}\}$


Factor with $\{D, \{A, C\}\}$

Therefore $(B \perp D|A, C)$

Similarly $(A \perp C|B, D)$

Basic Independencies

- As in Bayesian Networks, graph structure in a Markov network encodes a set of independence assumptions
- In a MN Probabilistic influence flows along the undirected paths in the graph and “*blocked*” if we condition on intervening nodes
 - i.e., we know their values
- We state this formally, next




*C blocks nodes
of A and B* 4


Active Path Definition

- Let H be a Markov network with nodes $\chi = \{X_1, \dots, X_n\}$,
- Let $X_1 - \dots - X_k$ be a path in H
- Let $Z \subseteq \chi$ be a set of observed variables
- A Path $X_1 - \dots - X_k$ is active given Z if none of the X_i is in Z

- Ex 1: If $Z = \{B\}$, path $A-D-C$ is active



- Ex 2: If $Z = C$, paths between nodes of A and C are inactive





- We can define separation in the graph when there is no active path, next

Separation and Global Independencies


1. Set of nodes Z separates sets X and Y denoted $sep_H(X; Y|Z)$ if there is no active path between any $X \in X$ and $Y \in Y$

- Ex 1: $Z = \{B, D\}$ separates A and C
 - i.e., $sep_H(A; C|B, D)$ or there is no active path between A and C
- Ex 2: C separates A and B



2. Global independencies associated with H are

- $I(H) = \{(X \perp Y|Z) : sep_H(X; Y|Z)\}$
- Independencies in $I(H)$ guaranteed to hold for every distribution P over H



$$I(H) = \{(A \perp C|B, D), (B \perp D|A, C)\}$$

Definition of Separation Leads to a disadvantage

- With a superset of Z , separation still holds
 - If $sep_H(X; Y|Z)$ then $sep_H(X; Y|Z')$ for any $Z' \supset Z$
- If separation is taken as definition of independencies, we restrict ability to encode non-monotonic independence relations
 - Non-monotonic reasoning is quite useful
 - E.g., intercausal reasoning with BNs
 - Two diseases are independent, but dependent given some common symptom
 - Such independence properties cannot be expressed as a Markov network

Factorization and Independencies

- Can show connection between independence properties implied by a Markov structure and factorizing a distribution over the graph
- Analogous to Bayesian Networks
 - Let G be a BN for a set of random variables χ and P be a distribution over χ .
 - If P factorizes according to G , i.e., product of CPDs, then G is an I-map of P
 - i.e., independencies $I(G) \subseteq I$

Formalizing independencies in MNs and distributions

- Gibbs Distribution

- A distribution P_Φ is a Gibbs distribution parameterized by a set of factors $\Phi = \{\phi_1(D_1), \dots, \phi_K(D_K)\}$
- If defined as follows

$$P_\Phi(X_1, \dots, X_n) = \frac{1}{Z} \tilde{P}(X_1, \dots, X_n)$$

where

$$\tilde{P}(X_1, \dots, X_n) = \prod_{i=1}^m \phi_i(D_i)$$

is an unnormalized measure and

$$Z = \sum_{X_1, \dots, X_n} \tilde{P}(X_1, \dots, X_n)$$
 is a normalizing constant

called the partition function

D_i are sets of random variables

Soundness of Separation Criterion

- Theorem 1 (from factorization to independencies):
 - Let P be distributed over $\chi = \{X_1, \dots, X_n\}$ and H a Markov structure over χ
 - If P is a Gibbs distribution that factorizes over H , (i.e., every D_i in H is a clique), then H is an I-map for P (i.e., every independency in H holds in P)
- Theorem 2: Hammersley-Clifford
(other direction: from independencies to factorization)
 - if H is an I-map for P then P factorizes over H
 - Holds only for positive distributions ($P > 0$ for every assignment)

Positive Distribution


- A distribution P is said to be positive if for all events $\alpha \in S$ such that $\alpha \neq \Phi$ we have that $P(\alpha) > 0$
- A non-positive distribution
- 16 possible values
 - Distribution P : 8 have value 1/8 rest are zero

X_1	X_2	X_3	X_4	$P(X)$
0	0	0	0	1/8
1	0	0	0	1/8
1	1	0	0	1/8
1	1	1	0	1/8
0	0	0	1	1/8
0	0	1	1	1/8
0	1	1	1	1/8
1	1	1	1	1/8

Rest 8 probs are 0

Non-positive distribution consistent with H

- Four binary random variables
- Global Independencies in graph H:
 - Consider $X_1 - X_2 - X_3 - X_4 - X_1$
 - implies $(X_1 \perp X_3 | X_2, X_4)$
- P also satisfies this
 - For the assignment $X_2=1, X_4=0$
 - $P(X_1=1 | X_2=1, X_4=0) = 1$
 - Rest are zero
 - Thus X_1 is independent of X_3




X_1	X_2	X_3	X_4	$P(\mathbf{X})$
0	0	0	0	1/8
1	0	0	0	1/8
1	1	0	0	1/8
1	1	1	0	1/8
0	0	0	1	1/8
0	0	1	1	1/8
0	1	1	1	1/8
1	1	1	1	1/8

Rest 8 probs are 0

- Global independencies hold \Rightarrow H is an I-map for P
- But P does not factorize according to H (Proof by contradiction)¹²

Graphical Model as Filter



$p(x)$ is allowed to pass through only if
It satisfies independencies in graph
This set is denoted DF or UF (for BN or MN)

- UI is set of distributions that are consistent with set of conditional independence statements read from the undirected graph using graph separation
- UF are set of distributions that can be expressed as factorization of the form

$$p(\mathbf{x}) = \frac{1}{Z} \prod_c \psi_c(\mathbf{x}_c)$$

- *Hammersley-Clifford theorem* states that UI and UF are identical

Independencies in Bayesian Networks

- Bayesian networks have two types of independencies
 - Local independencies
 - Each node is independent of its non-descendants given its parents
 - Global independencies
 - Induced by d-separation
- These two sets of independencies are equivalent
 - One implies the other

Three Independencies of Markov Networks

- Two local independencies
 - Pairwise $I_p(H)$
 - Markov Blanket $I_l(H)$
- Global independency $I(H)$
 - Identify three sets of nodes A , B and C
 - To test conditional independence property
$$A \perp B | C$$
 - Consider all possible paths from nodes in set A to nodes in set B
 - If all such paths pass through one or more nodes in C then path is blocked and independence holds


Pairwise Independency $I_p(H)$

- Weakest independency
 - Whenever two variables are directly connected
 - Potential of being **correlated** not mediated by other variables
 - When two variables are not directly linked
 - There must be a way of rendering them **conditionally independent**
- If H is a Markov network its *pairwise independencies* are

$$I_p(H) = \{(X \perp Y | \chi - \{X, Y\}) : X - Y \notin H\}$$

- Example


$$I_p(H) = \{(A \perp C | B, D), (B \perp D | A, C)\}$$



When X, Y are **not** directly connected i.e., $X - Y \notin H$, they are **independent** given all other variables

Markov Blanket Independency $I_I(H)$

- Analogous to local independencies in Bayesian networks
 - We can block all influences by conditioning on its immediate neighbors
 - Node is conditionally independent of all nodes given its immediate neighbors
- For graph H the Markov blanket of X in H is the set of neighbors of X in H
- Local independencies associated are
 - $I_I(H) = \{(X \perp \chi - \{X\} - MB_H(X) \mid MB_H(X)) : X \in \chi\}$



Relationship between Markov properties

- Three independencies of network structure H
- $I_p(H)$ is strictly weaker than $I_l(H)$ is strictly weaker than $I(H)$
- For positive distributions all three are equivalent

Separation in Markov Networks

- Markov network encodes a set of conditional independencies

- Probabilistic influence flows

- in undirected paths

- Blocked if we condition on intervening nodes

- Every path from any node in A to B passes through C

- No explaining away


- Testing for independence simpler than in directed graphs

- Alternative view


- Remove all nodes in set C together with all their connecting links

- If no paths from A to B then conditional independence holds

- Markov blanket



- A node is conditionally independent of all other nodes conditioned only on its neighbors



Factorization Properties

- Factorization rule corresponds to conditional independence test
- Notion of locality needed
- Consider two nodes x_i and x_j not connected by a link
 - They are conditionally independent given all other nodes in graph
 - Because there is no direct path between them and
 - All other paths pass through nodes that are observed and hence those paths are blocked
 - Expressed as

$$p(x_i, x_j \mid \mathbf{X} \setminus \{i, j\}) = p(x_i \mid \mathbf{X} \setminus \{i, j\}) p(x_j \mid \mathbf{X} \setminus \{i, j\})$$
 - Where $\mathbf{X} \setminus \{i, j\}$ denotes set \mathbf{x} of all variables with x_i and x_j removed
- For conditional independence to hold
 - factorization is such that x_i and x_j do not appear in the same factor
 - No path between them other than going through others
 - leads to graph concept of clique

