

# Unsupervised Learning

Sargur Srihari

[srihari@cedar.buffalo.edu](mailto:srihari@cedar.buffalo.edu)

# Topics in Mixture Models and EM

- Unsupervised Learning and Mixture Models
  1. K-means algorithm for finding clusters in a data set
  2. Latent variable view of mixture distributions
  3. General technique for finding m.l. estimators in latent variable models
  4. EM Algorithm
  5. Infinite Mixture Models

# Unsupervised vs Supervised

- Unsupervised algorithms experience only “features” but not a supervision signal
- Distinction between supervised and unsupervised algorithms is not formally and rigidly defined
  - Because there is no objective test for distinguishing whether a value is a feature or a target provided by a supervisor
- Informally unsupervised learning refers to attempts to extract information from a distribution that do not require human labor to annotate examples


# Unsupervised Algorithms

- Density estimation

- Learning to draw samples from a distribution
- Learning to denoise data from a distribution




- Clustering data into related groups




- Finding the best representation

1. Lower dimensional representations
2. Finding a manifold that the data lies near
3. Sparse representations
4. Independent representations



- Autoencoders



# Intelligence as a cake

1. The targets for supervised learning contain *far less* information than input data
2. Reinforcement Learning reward contains *even less*
3. Unsupervised learning gives us an essentially unlimited supply of information about the world



Unsupervised learning would be the cake



Supervised learning would be the icing on the cake

RL would be the cherry on the cake

# Simple representation learning

## 1. Principal components analysis

Original Data  $\mathbf{x}$




Transformed Data  $\mathbf{z} = \mathbf{x}^T \mathbf{W}$

Aligns axes along direction of greatest variance


## 2. $k$ - means clustering

## 3. Mixture models and EM


Original (Synthetic) Data



K-means Clustering




GMM & EM Clustering



# Autoencoder


- VAE is a *generative model*
  - Generates samples that look like training data



- Due to random variable between input-output it cannot be trained using backprop
- Backprop through parameters of latent distribution
  - Called reparameterization trick


$$N(\mu, \Sigma) = \mu + \Sigma N(0, I)$$

Where  $\Sigma$  is diagonal



# Modeling complex distributions

- Complex distribution  $p(x)$  of observed variable  $x$




$x$

- Can be expressed in terms of a more tractable joint distribution over observed and latent variables  $p(x, z)$ 
  - Latent variable  $z$  with three values can model this distribution
- Distribution of  $x$  alone obtained by marginalization  $p(x) = \sum_z p(x, z)$
- Latent variables allow complicated distributions to be formed from simpler components
  - Gaussian mixtures have latent variables  $z$  that are discrete
  - Also called Finite Mixture Models


# Gaussian Mixture Model (GMM)

- Linear superposition of Gaussian components
  - Two Gaussians
  - Three Gaussians

$z$  has 2 values



$z$  has 3 values (data: petal width in Iris)



Since

$$p(x) = \sum_z p(x, z) = \sum_z p(z)p(x|z)$$

We can write (for the mixture of two Gaussians):

$$p(x) = p(z=1)p(x|z=1) + p(z=2)p(x|z=2)$$

# Mixture Model As Unsupervised Learning


- Probabilistic model representing sub-populations within a population
  - Without requiring that the sub-population of the data items be identified (supervised)
- Constructing such models is called unsupervised learning or clustering
- Can be applied to discrete distributions as well

# Bernoulli Mixture Model


- Clustering handwritten Digit Data ( $560 \times 420$  pixels)
  - Mixture Model for digits 0-4 with  $K=12$

- 12 {
- Identifies three types of 0s (elongated, circular and tilted)
  - two 1s (straight and tilted)
  - two 2s (loopy and bird)
  - two 3s (forward and backward leaning)
  - three 4s (vertical bars)

Superimposed data of 12 components



Examples of 12 components



# Role of Mixture Models

- Mixture models provide:
  1. Framework for building complex probability distributions
    - Complex distribution expressed in terms of tractable joint distribution of observed and latent variables
      - Distribution of observed variables: by marginalization
  2. A method for clustering data
    - Unsupervised learning