

PGM Regression

Sargur Srihari

srihari@cedar.buffalo.edu


Topics

1. What are graphical models (or PGMs)
2. Directed and Undirected Graphical Models
3. Joint and Conditional Probability Distributions
4. Probabilistic Queries
- 5. Regression expressed as PGMs**
6. PGMs as Generative Models and Sampling
7. PGM for Discrete Distributions

Example: Bayesian Polynomial Regression

- Directed graph to describe probability distribution of scalar x
- The polynomial regression problem

$$y(x, \mathbf{w}) = w_0 + w_1x + w_2x^2 + \dots + w_Mx^M$$



- Probabilistic formulation with Random variables:
 - Vector of polynomial coefficients \mathbf{w}
 - Observed data $\mathbf{t} = (t_1, \dots, t_N)^T$ where $t_n = y(x_n, \mathbf{w}) + \varepsilon$ is a noise corrupted version of x_n
- Focusing on random variables, joint distribution is given by

From product rule $p(\mathbf{t}, \mathbf{w}) = p(\mathbf{w}) p(\mathbf{t} | \mathbf{w})$

$$p(\mathbf{t}, \mathbf{w}) = p(\mathbf{w}) \prod_{n=1}^N p(t_n | \mathbf{w})$$

prior

Conditional distributions



Since samples are independent

Graph for Polynomial Regression

- Joint distribution is given by

$$p(t, \mathbf{w}) = p(\mathbf{w}) \prod_{n=1}^N p(t_n | \mathbf{w})$$

- It is to be represented by a graphical model
 - Observing rhs introduce a node for each variable
 - Introduce associations
 - A link for each conditional distribution
- Equivalent *plate* representation



Parameters in Bayesian Poly Regression

- Random variables:
 - Vector of polynomial coefficients w
 - Observed data $t = (t_1, \dots, t_N)^T$ where t_n is a noise corrupted version of x_n
 - Additionally model contains:
 - Input data $x = (x_1, \dots, x_N)^T$
 - Noise variance σ^2
 - Hyper-parameter α , which is the precision (inverse variance) of Gaussian prior
- Parameters of model rather than random variables
- Focusing on random variables, joint distribution is given by


$$p(t, w) = \underbrace{p(w)}_{\text{Gaussian prior with precision } \alpha} \prod_{n=1}^N \underbrace{p(t_n | w)}_{\text{Conditional distributions with variance } \sigma^2}$$

With Deterministic Parameters

- Sometimes useful to explicitly show parameters of model
- Same model with deterministic parameters


$$p(t, w | x, \alpha, \sigma^2) = p(w | \alpha) \prod_{n=1}^N p(t_n | w, x_n, \sigma^2)$$

- Random variables are denoted by open circles
- Deterministic parameters by smaller (tiny) solid circles




Observed and Latent Variables

- In ML some random variables are set to specific observed values
 - Observed variable nodes are shaded
- Variable w is not observed
 - It is *latent* or *hidden*
- Nodes t_n are shaded
 - to indicate corresponding variables are set to observed values



Including predictions

- Polynomial Regression model including input and predicted values
- Need to write out longer probability expression including
 - new input value \hat{x} and model prediction \hat{t}
- Joint distribution of all variables in this model



$$p(\hat{t}, t, w | \hat{x}, x, \alpha, \sigma^2) = p(w | \alpha) \prod_{n=1}^N p(t_n | w, x_n, \sigma^2) p(\hat{t} | \hat{x}, w, \sigma^2)$$

- Predictive distribution

$$p(\hat{t} | \hat{x}, x, \alpha, \sigma^2) \propto \int p(\hat{t}, t, w | \hat{x}, x, \alpha, \sigma^2) dw$$

