

Probability-Foundations

Sargur Srihari

srihari@cedar.buffalo.edu

Topics

- Random Variables, Joint and Conditional Distributions
- Independence and Conditional Independence
- Answering Queries
 - Probabilistic queries
 - What is the probability of this event?
 - MAP queries, Marginal MAP queries
 - What is the most probable setting of these variables?
 - Most Possible Explanation queries
 - What is the most probable setting of these variables given the evidence

Random Variable

- We have a population of students
 - We want to reason about their grades
 - Consider just one random variable: *Grade*
 - $P(\text{Grade})$ associates a probability with each outcome
 $\text{Val}(\text{Grade}) = \{ A, B, C \}$
- If $k = |\text{Val}(X)|$ then $\sum_{i=1}^k P(X = x^i) = 1$
 - k is no of elements in $\text{Val}(X)$
 - x^i are the elements in $\text{Val}(X)$
 - Distribution with $k > 2$ is a *multinomial* distribution
 - If $\text{Val}(X) = \{ \text{false}, \text{true} \}$, or $k = 2$, it is a *Bernoulli* distribution
- $P(X)$ is known as the marginal distribution of X

Joint Distribution

- We are interested in questions involving several random variables
 - Need to consider joint distributions
 - Over a set $\chi = \{X_1, \dots, X_n\}$ denoted by $P(X_1, \dots, X_n)$
 - We use ξ to refer to a full assignment to variables χ , i.e. $\xi \in \text{Val}(\chi)$
- Example of joint distribution of two variables:
 - $P(G, I)$: *Intelligence (I) and Grade(G)*
 - And marginal distributions
 - $P(G), P(I)$

$P(G, I)$	Intelligence		$P(G)$
	low	high	
Grade	A	0.07 0.18	0.25
	B	0.28 0.09	0.37
	C	0.35 0.03	0.38
$P(I)$		0.7 0.3	1

Conditional Probability

- $P(\textit{Intelligence}|\textit{Grade}=A)$ describes the distribution over events describable by *Intelligence* given knowledge that student's grade is A
- It is not the same as the marginal distribution

$P(G,I)$		<i>Intelligence</i>		
		<i>low</i>	<i>high</i>	
<i>Grade</i>	A	0.07	0.18	0.25
	B	0.28	0.09	0.37
	C	0.35	0.03	0.38
		0.7	0.3	1

$$P(\textit{Intelligence}=\textit{high})=0.3$$

$$P(\textit{Intelligence}=\textit{high}|\textit{Grade}=A)$$

$$=0.18/0.25$$

$$=0.72$$

Marginal and
Conditional
are very different

Independent Random Variables

- We expect $P(\alpha|\beta)$ to be different from $P(\alpha)$
 - i.e., β is true changes our probability over α
- Sometimes equality can occur, i.e., $P(\alpha|\beta)=P(\alpha)$
 - i.e., learning that β occurs did not change our probability of α
 - We say event α is independent of event β denoted $P \rightarrow (\alpha \perp \beta)$ if $P(\alpha|\beta)=P(\alpha)$ or if $P(\beta)=0$
- A distribution P satisfies $(\alpha \perp \beta)$ if and only if $P(\alpha \wedge \beta)=P(\alpha)P(\beta)$


Conditional Independence

- While independence is a useful property, we don't often encounter two independent events
- A more common situation is when two events are independent given an additional event
 - Reason about student job offer at *Google* or *Apple*
 - These two are not independent
 - If student offered at *Google* then probability of *Apple* is higher
 - If both based on GPA and we know the GPA to be *A*
 - Then the student being offered at *Apple* does not change probability of being offered at *Google*
 - $P(\text{Apple} | \text{Google}, \text{Grade } A) = P(\text{Apple} | \text{Grade } A)$
 - i.e., *Apple* is conditionally independent of *Google* given *Grade A*

Querying Joint Probability Distributions

Evaluating a Joint Probability

Graph & CPDs



Chain rule for BN:

$$P(D, I, G, S, L) = P(D)P(I)P(G|D, I)P(S|I)P(L|G)$$

$$P(i^1, d^0, g^2, s^1, l^0) = P(i^1)P(d^0)P(g^2 | i^1, d^0)P(s^1 | i^1)P(l^0 | g^2)$$

$$= 0.3 \cdot 0.6 \cdot 0.08 \cdot 0.8 \cdot 0.4 = 0.004608$$

$Val(I) = \{i^0 = \text{low intelligence}, i^1 = \text{high intelligence}\}$
 $Val(D) = \{d^0 = \text{easy}, d^1 = \text{hard}\}$
 $Val(G) = \{g^1 = A, g^2 = B, g^3 = C\}$
 $Val(S) = \{s^0 = \text{low}, s^1 = \text{high}\}$
 $Val(L) = \{l^0 = \text{weak}, l^1 = \text{strong}\}$

$P(\text{high intelligence, easy course, grade=B, high SAT, weak letter}) = \text{very low}$



Query Types

1. Probability Queries

- Given L and S give probability distribution of I

2. MAP Queries

- Maximum a posteriori probability
- Also called MPE (*Most Probable Explanation*)
 - What is the most likely setting of D, I, G, S, L
 - Instead of probabilities we want values for the variables
- Marginal MAP Queries
 - When some variables are known

Probability Queries

- Most common type of query is a probability query
- Query has two parts
 - *Evidence*: a subset E of variables and their instantiation e
 - *Query Variables*: a subset Y of random variables in network
- **Inference Task**: $P(Y|E=e)$
 - *Posterior probability distribution* over values y of Y
 - *Conditioned* on the fact $E=e$
 - Can be viewed as Marginal over Y in distribution we obtain by conditioning on e
- **Marginal Probability Estimation**

$$P(Y = y_i | E = e) = \frac{P(Y = y_i, E = e)}{P(E = e)}$$

MAP Queries (Most Probable Explanation)

- Finding a high probability assignment to some subset of variables
- Most likely assignment to all non-evidence variables $W = \mathcal{X} - Y$

$$MAP(W | e) = \arg \max_w P(w, e)$$

Value of w for which $P(w, e)$ is maximum

- Difference from probability query
 - Instead of a probability we get the most likely value for all remaining variables

Example of MAP Queries

Medical Diagnosis Problem

Disease (A) causes Symptom (B)

Two possible Diseases

$$Val(A) = \{ a^0 = Mono, a^1 = Flu \}$$

Two possible Symptoms

$$Val(B) = \{ b^0 = Headache, b^1 = Fever \}$$

Marginal MAP Query

Q₁: Most likely disease $P(A)$?

A₁: Flu (Trivially determined for root node)

MAP Query

Q₂: Most likely disease and symptom $P(A, B)$?

A₂:

Marginal MAP Query

Q₃: Most likely symptom $P(B)$?

A₃:


$P(A)$

a^0

a^1

0.4

0.6



Disease

A

B

Symptom


$P(B|A)$

$P(B A)$	b^0	b^1
a^0	0.1	0.9
a^1	0.5	0.5

Evaluating Answers to Queries

$$P(A)$$

a^0	a^1
0.4	0.6



$$MAP(A) = \arg \max_a P(A = a^1) \quad A_1: \text{Flu}$$

$$MAP(A, B) = \arg \max_{a, b} P(A, B) = \arg \max_{a, b} P(A)P(B|A)$$

$$= \arg \max_{a, b} \{0.04, 0.36, 0.3, 0.3\} = a^0, b^1$$

A_2 : Mono and Fever

Note that individually most likely value a^1 is not in the most likely joint assignment

$P(B A)$	b^0	b^1
a^0	0.1	0.9
a^1	0.5	0.5

Marginal MAP Query

In MAP we looked for highest joint probability assignment of *Disease* and *Symptom*

Can look for most likely assignment of *Disease* variable only

Query is not all remaining variables but a subset of them

Y is query, evidence is $E=e$


What is the most probable explanation of answer to query *Y* ?

Task is to find most likely assignment to *Y*:

$$MAP(Y|e) = \arg \max P(Y|e)$$


$$MAP(Y|e) = \arg \max_Y \sum_Z P(Y,Z|e)$$

If $Z = \chi - Y - E$



<i>A</i>	b^0	b^1
a^0	0.1	0.9
a^1	0.5	0.5

Example of Marginal MAP Query



$P(B A)$	b^0	b^1
a^0	0.1	0.9
a^1	0.5	0.5

$$\begin{aligned}
 MAP(B) &= \arg \max_b P(B) = \arg \max_b \sum_A P(A, B) \\
 &= \arg \max_b \{0.34, 0.66\} = b^1
 \end{aligned}$$

A3: Fever

$P(A, B)$	b^0	b^1
a^0	0.04	0.36
a^1	0.3	0.3

Non-monotonic Marginal MAP

- Marginal MAPs are not monotonic
- Most likely assignment $MAP(Y_1|e)$ might be completely different from assignment to Y_1 in $MAP(\{Y_1, Y_2\}|e)$
 - Q_1 : Most likely disease $P(A)$?
 - A_1 : Flu
 - Q_2 : Most likely disease and symptom $P(A, B)$?
 - A_2 : Mono and Fever
- Thus we cannot use a MAP query to give a correct answer to a marginal map query

Marginal MAP more Complex than MAP

- Contains both summations (like in probability queries) and maximizations (like in MAP queries)

$$\begin{aligned} MAP(B) &= \arg \max_b P(B) = \arg \max_b \sum_A P(A, B) \\ &= \arg \max_b \{0.34, 0.66\} = b^1 \end{aligned}$$