

# I-Maps: Graphs and Distributions

Sargur Srihari  
srihari@cedar.buffalo.edu

# Topics

- I-Maps
- I-Map to Factorization
- Factorization to I-Map
- Perfect Map

# Graphs and Distributions

- Relating two concepts:
  - Independencies in distributions
  - Independencies in graphs
- I-Map is a relationship between the two

# Independencies in a Distribution

- Let  $P$  be a distribution over  $X$
- Define  $I(P)$  to be the set of conditional independence assertions of the form  $(X \perp Y | Z)$  that hold in  $P$
- **Example: Joint distribution  $P(X, Y)$  given by table**

$X$	$Y$	$P(X, Y)$
$x^0$	$y^0$	0.08
$x^0$	$y^1$	0.32
$x^1$	$y^0$	0.12
$x^1$	$y^1$	0.48

$X$  and  $Y$  are independent in  $P$ , e.g.,


$$P(x^1) = P(x^1, y^0) + P(x^1, y^1) = 0.12 + 0.48 = 0.6$$

$$P(y^1) = 0.32 + 0.48 = 0.8$$

$$P(x^1, y^1) = 0.48 = 0.6 \times 0.8$$

Thus  $(X \perp Y | \phi) \in I(P)$

# Independencies in a Graph



- Graph  $G$  with CPDs is equivalent to a set of independence assertions

$$P(D, I, G, S, L) = P(D)P(I)P(G | D, I)P(S | I)P(L | G)$$

- Local Conditional Independence Assertions** (starting from leaf nodes):

$$I(G) = \{(L \perp I, D, S | G), \quad L \text{ is conditionally independent of all other nodes given parent } G$$

$$(S \perp D, G, L | I), \quad S \text{ is conditionally independent of all other nodes given parent } I$$

$$(G \perp S | D, I), \quad \text{Even given parents, } G \text{ is NOT independent of descendant } L$$

$$(I \perp D | \phi), \quad \text{Nodes with no parents are marginally independent}$$


$$(D \perp I, S | \phi)\} \quad D \text{ is independent of non-descendants } I \text{ and } S$$

- Parents of a variable shield it from probabilistic influence
  - Once value of parents known, no influence of ancestors
- Information about descendants can change beliefs about a node


# Definition of I-MAP

- Let  $\mathcal{K}$  be a graph associated with a set of independencies  $I(\mathcal{K})$
- Let  $P$  be a probability distribution with a set of independencies  $I(P)$
- Then  $\mathcal{K}$  is an  $I$ -map of  $P$  if  $I(\mathcal{K}) \subseteq I(P)$ 
  - From direction of inclusion
    - Distribution has more independencies than graph
    - Graph doesn't mislead in independencies in  $P$ 
      - Any independence that  $G$  asserts must also hold in  $P$


# I-Map Examples: $G$ and $P$



$G_0$  encodes  $X \perp Y$  or  $I(G_0) = \{X \perp Y\}$



$G_1$  encodes no Independence or  $I(G_1) = \{\emptyset\}$



$G_2$  encodes no Independence  $I(G_2) = \{\emptyset\}$

$P$ :  $X$  and  $Y$  are independent

$X$	$Y$	$P(X, Y)$
$x^0$	$y^0$	0.08
$x^0$	$y^1$	0.32
$x^1$	$y^0$	0.12
$x^1$	$y^1$	0.48

$G_0$  is an I-map of  $P$   
 $G_1$  is an I-map of  $P$   
 $G_2$  is an I-map of  $P$

$P$ :  $X$  and  $Y$  are **not** independent, e.g.,

$X$	$Y$	$P(X, Y)$
$x^0$	$y^0$	0.4
$x^0$	$y^1$	0.3
$x^1$	$y^0$	0.2
$x^1$	$y^1$	0.1

$P(x^1, y^1) = 0.1$   
 $P(x^1) = 0.3$   $P(y^1) = 0.4$

Thus

$$\boxed{(X \perp Y) \notin I(P)}$$

$G_0$  is not an I-map of  $P$   
 $G_1$  is an I-map of  $P$   
 $G_2$  is an I-map of  $P$

If  $G$  is an I-map of  $P$  then it captures some of the independences, not all

# I-map to Factorization

- A Bayesian network  $G$  encodes a set of conditional independence assumptions  $I(G)$
- Every distribution  $P$  for which  $G$  is an I-map should satisfy these assumptions
  - Every element of  $I(G)$  should be in  $I(P)$
- This is the key property to allowing a compact representation

# I-map to Factorization

- Consider Joint distribution  $P(I, D, G, L, S)$ 
  - From chain rule of probability
 
$$P(I, D, G, L, S) = P(I)P(D|I)P(G|I, D)P(L|I, D, G)P(S|I, D, G, L)$$
  - Relies on no assumptions, Also not very helpful
    - Last factor requires evaluation of 24 conditional probabilities
  - Assume  $G_{\text{student}}$  is an I-map
    - Apply conditional independence assumptions induced from the graph

$D \perp I \in I(P)$  therefore  $P(D|I) = P(D)$


$(L \perp I, D) \in I(P)$  therefore  $P(L|I, D, G) = P(L|G)$

– Thus we get

$$P(D, I, G, S, L) = P(D)P(I)P(G|D, I)P(S|I)P(L|G)$$

- Which is a factorization into local probability models

– Thus we can go from graphs to factorization of  $P$




# Factorization to I-map

- We have seen that we can go from the independences encoded in  $G$ , i.e.,  $I(G)$ , to Factorization of  $P$
- Conversely, Factorization according to  $G$  implies associated conditional independences
  - If  $P$  factorizes according to  $G$  then  $G$  is an I-map for  $P$
  - Need to show that if  $P$  factorizes according to  $G$  then  $I(G)$  holds in  $P$
  - Proof by example

# Example that independences in $G$ hold in $P$

- $P$  is defined by set of CPDs
- Consider independences for  $S$  in  $G$ , i.e.,  

$$P(S \perp D, G, L | I)$$



- Starting from factorization induced by graph


$$P(D, I, G, S, L) = P(D)P(I)P(G | D, I)P(S | I)P(L | G)$$

- Can show that  $P(S | I, D, G, L) = P(S | I)$
- Which is what we had assumed for  $P$

# Perfect Map


- I-map

- All independencies in  $I(G)$  present in  $I(P)$
- Trivial case: all nodes interconnected



- D-Map

- All independencies in  $I(P)$  present in  $I(G)$
- Trivial case: all nodes disconnected



- Perfect map

- Both an I-map and a  $D$ -map
- Interestingly not all distributions  $P$  over a given set of variables can be represented as a perfect map

- Venn Diagram where  $D$  is set of distributions that can be represented as a perfect map

