


Local Probabilistic Models: Tabular CPDs

Sargur Srihari

srihari@cedar.buffalo.edu

Local Probabilistic Models

- Bayesian Networks capture *global properties* of independence of variables



$$I(G) = \{(D \perp I \mid \phi), (G \perp S \mid D, I), (S \perp D, G, L \mid I), (D \perp I, S \mid \phi), (L \perp I, D, S \mid G)\}$$

- Properties of independence allow us to:
 - factorize high-dimensional joint distribution into product of lower-dimensional CPDs (or factors)

$$P(D, I, G, S, L) = P(D)P(I)P(G \mid D, I)P(S \mid I)P(L \mid G)$$

- Next: exploit additional regularities in CPDs

Local Probabilistic Model Types




TOC for Local Probabilistic Models

1. Tabular CPDs
2. Deterministic CPDs
3. Context-Specific CPDs
 - (1) Tree CPD (Printer Diagnosis), (2) Rule CPD
4. Independence of Causal Influence
 - (1) Noisy-OR, (2) Generalized Linear Models
5. Continuous Variables: Robotics
 - Hybrid Models: Thermostat
6. Conditional BNs: Computer Network

Tabular CPDs

- When Random Variables are discrete-valued



- Encode $P(X|pa_X)$ as a table

- Contains an entry for each assignment to X & pa_X
- Proper CPD requires all non-negative values and


$$\sum_{x \in Val(x)} P(x | pa_X) = 1$$

- Inference algorithms can use table CPDs in a natural way

- Leads to perception that table CPDs are inherent to BNs, but.....


Disadvantages of Tabular CPDs

- Continuous case
 - Cannot store each conditional probability in a table
 - Domain of random variable is infinite
- Discrete case
 - Parameters: exponential with no. of parents:
 - for X with n binary parents need 2^n values for $P(X|pa_X)$



Unwieldiness of CPTs

- Tabular representation becomes rapidly large as the no. of parents grows
- It is a serious one in many settings
 - If *Fever* is caused by 10 diseases, we need to ask expert to answer 1,024 questions— tiresome!



- Regularity among CPDs is not exploited
 - When D_1 true, *Fever* is certain irrespective of others

Solution: Different viewpoint

- A CPD needs to specify a conditional probability $P(x|\text{pa}_X)$ for every assignment of values pa_X and x but does not have to do so by listing each such value explicitly
- View the CPDs not as tables listing all conditional probabilities but as functions that given pa_X and x return the conditional probability $P(x|\text{pa}_X)$