

Local Probabilistic Models: Deterministic CPDs

Sargur Srihari

srihari@cedar.buffalo.edu

Topics

- **Local Probabilistic Models**
 1. Tabular CPDs
 - 2. Deterministic CPDs**
 3. Context-Specific CPDs
 - (1) Tree CPD (Printer Diagnosis), (2) Rule CPD
 4. Independence of Causal Influence
 - (1) Noisy-OR, (2) Generalized Linear Models
 5. Continuous Variables: Robotics
 - Hybrid Models: Thermostat
 6. Conditional BNs: Computer Network

Deterministic CPDs

- Simplest non-tabular CPD
- Child X : a deterministic function of parents pa_X ,
 - i.e., there is a function f such that
 - $f: \text{Val}(\text{pa}_X) \rightarrow \text{Val}(X)$



$$P(x | \text{pa}_X) = \begin{cases} 1 & x = f(\text{pa}_X) \\ 0 & \text{otherwise} \end{cases}$$

- Two examples
 - Ex: All are binary-valued variables: X is “or” of parents:

$$f: P(X|Y,Z) = Y \vee Z$$
 - Ex: Continuous domain: We want to assert in $P(X|Y,Z)$ that X is sum of parent values

Ex of Deterministic CPD: Modeling a car

- **Tires** T_1, T_2, T_3, T_4 $Var(T_i) = \{\text{flat}, \text{inflated}\}$
 - **Two effect variables:** Steering, Ride
- Instead of effects having as parents all T_i s, have them depend on single variable T
 - T : deterministic or of its parents
- **Advantages** $T = T_1 \vee T_2 \vee T_3 \vee T_4$
 - **Reduced indegree (8 vs 2)**
 - Each effect has 1 instead 4 parents
 - If there are more dependencies, considerable savings



Deterministic CPDs & Independencies


- Determining Independencies in a BN are slightly different with deterministic CPDs
- Recall that conditional independence is a numeric property

$$A \perp B | C \iff P(A, B | C) = P(A | C) P(B | C)$$


- Although defined using equality of probabilities
graph structure allows us to deduce some independencies without looking at the numbers
- Need to modify *D-separation* for determinism

Ex: Modifying D-separation

- If C is a deterministic function of A and B , what new independencies exist?
 - If A and B are known, C is known, so D and E are independent: $(D \perp E \mid A, B)$
 - Not necessarily true if C were not deterministic
 - If C is not known $D \not\perp E$



More Complex Example with Deterministic CPDs



- C is exclusive-or of A and B

$$C = A \oplus B$$

- If B and C are known, then A is known
 - Therefore D and E are independent: $(D \perp E \mid B, C)$
 - This would not be true if C were nondeterministic