

Maximum-Likelihood and Bayesian Parameter Estimation

- Sufficient Statistics
- Common Probability Distributions
- Problems of Dimensionality

Sufficient Statistics

- Any function s of the samples is a statistic
- A *sufficient statistic* is a (possibly vector valued) function s of samples D that contains all information relevant to estimating some parameter θ

Factorization Theorem:

A statistic s is sufficient for θ iff the probability $P(D/ \theta)$ can be written as the product

$$P(D/ \theta) = g(s, \theta) h(D)$$

For some functions $g(. , .)$ and $h(. , .)$

Sufficient Statistics and Gaussian

$$p(x | \theta) \sim N(\theta, \Sigma)$$







Unknown mean

$$\begin{aligned} p(D | \theta) &= \prod_{k=1}^n \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp\left[-\frac{1}{2} (x_k - \theta)^t \Sigma^{-1} (x_k - \theta)\right] \\ &= \exp\left[-\frac{n}{2} \theta^t \Sigma^{-1} \theta + \theta^t \Sigma^{-1} \left(\sum_{k=1}^n x_k\right)\right] \times \frac{1}{(2\pi)^{nd/2} |\Sigma|^{n/2}} \exp\left[-\frac{1}{2} \sum_{k=1}^n x_k^t \Sigma^{-1} x_k\right] \end{aligned}$$






Isolates dependence on θ
in the first term

Sufficient Statistic

Common Exponential Distributions and their Sufficient Statistics

| Name | Distribution | Domain | | s | $[g(s, \theta)]^{1/n}$ |
|---------------------|--|-----------------------------------|--|---|--|
| Normal | $p(x \theta) = \sqrt{\frac{\theta_2}{2\pi}} e^{-(1/2)\theta_2(x-\theta_1)^2}$ | $\theta_2 > 0$ |  | $\begin{bmatrix} \frac{1}{n} \sum_{k=1}^n x_k \\ \frac{1}{n} \sum_{k=1}^n x_k^2 \end{bmatrix}$ | $\sqrt{\theta_2} e^{-\frac{1}{2}\theta_2(s_2 - 2\theta_1 s_1 + \theta_1^2)}$ |
| Multivariate Normal | $p(\mathbf{x} \theta) = \frac{ \Theta_2 ^{1/2}}{(2\pi)^{d/2}} e^{-(1/2)(\mathbf{x}-\theta_1)^t \Theta_2 (\mathbf{x}-\theta_1)}$ | Θ_2 positive definite |  | $\begin{bmatrix} \frac{1}{n} \sum_{k=1}^n \mathbf{X}_k \\ \frac{1}{n} \sum_{k=1}^n \mathbf{X}_k \mathbf{X}_k^t \end{bmatrix}$ | $ \Theta_2 ^{1/2} e^{-\frac{1}{2}[\text{tr}\Theta_2 s_2 - 2\theta_1^t \Theta_2 s_1 + \theta_1^t \Theta_2 \theta_1]}$ |
| Exponential | $p(x \theta) = \begin{cases} \theta e^{-\theta x} & x \geq 0 \\ 0 & \text{otherwise} \end{cases}$ | $\theta > 0$ |  | $\frac{1}{n} \sum_{k=1}^n x_k$ | $\theta e^{-\theta s}$ |
| Rayleigh | $p(x \theta) = \begin{cases} 2\theta x e^{-\theta x^2} & x \geq 0 \\ 0 & \text{otherwise} \end{cases}$ | $\theta > 0$ |  | $\frac{1}{n} \sum_{k=1}^n x_k^2$ | $\theta e^{-\theta s}$ |
| Maxwell | $p(x \theta) = \begin{cases} \frac{4}{\sqrt{\pi}} \theta^{3/2} x^2 e^{-\theta x^2} & x \geq 0 \\ 0 & \text{otherwise} \end{cases}$ | $\theta > 0$ |  | $\frac{1}{n} \sum_{k=1}^n x_k^2$ | $\theta^{3/2} e^{-\theta s}$ |
| Gamma | $p(x \theta) = \begin{cases} \frac{\theta_2^{\theta_1+1}}{\Gamma(\theta_1+1)} x^{\theta_1} e^{-\theta_2 x} & x \geq 0 \\ 0 & \text{otherwise} \end{cases}$ | $\theta_1 > -1$ $\theta_2 > 0$ |  | $\begin{bmatrix} \left(\prod_{k=1}^n x_k \right)^{1/n} \\ \frac{1}{n} \sum_{k=1}^n x_k \end{bmatrix}$ | $\frac{\theta_2^{\theta_1+1}}{\Gamma(\theta_1+1)} s^{\theta_1} e^{-\theta_2 s}$ |

Common Exponential Distributions and their Sufficient Statistics

| Name | Distribution | Domain | s | $[g(s, \theta)]^{1/n}$ | |
|-------------|--|---|---|---|--|
| Beta | $p(x \theta) = \begin{cases} \frac{\Gamma(\theta_1+\theta_2+2)}{\Gamma(\theta_1+1)\Gamma(\theta_2+1)} x^{\theta_1} (1-x)^{\theta_2} & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$ | $\theta_1 > -1$ $\theta_2 > -1$ |  | $\left[\begin{matrix} \left(\prod_{k=1}^n x_k \right)^{1/n} \\ \left(\prod_{k=1}^n (1-x_k) \right)^{1/n} \end{matrix} \right]$ | $\frac{\Gamma(\theta_1+\theta_2+2)}{\Gamma(\theta_1+1)\Gamma(\theta_2+1)} s^{\theta_1} s^{\theta_2}$ |
| Poisson | $P(x \theta) = \frac{\theta^x}{x!} e^{-\theta} \quad x = 0, 1, 2, \dots$ | $\theta > 0$ |  | $\frac{1}{n} \sum_{k=1}^n x_k$ | $\theta^s e^{-\theta}$ |
| Bernoulli | $P(x \theta) = \theta^x (1-\theta)^{1-x} \quad x = 0, 1$ | $0 < \theta < 1$ |  | $\frac{1}{n} \sum_{k=1}^n x_k$ | $\theta^s (1-\theta)^{1-s}$ |
| Binomial | $P(x \theta) = \frac{m!}{x!(m-x)!} \theta^x (1-\theta)^{m-x} \quad x = 0, 1, \dots, m$ | $0 < \theta < 1$ |  | $\frac{1}{n} \sum_{k=1}^n x_k$ | $\theta^s (1-\theta)^{m-s}$ |
| Multinomial | $P(\mathbf{x} \theta) = \frac{m! \prod_{i=1}^d \theta_i^{x_i}}{\prod_{i=1}^d x_i!} \quad \begin{matrix} x_i = 0, 1, \dots, m \\ \sum_{i=1}^d x_i = m \end{matrix}$ | $0 < \theta_i < 1$ $\sum_{i=1}^d \theta_i = 1$ |  | $\frac{1}{n} \sum_{k=1}^n \mathbf{X}_k$ | $\prod_{i=1}^d \theta_i^{s_i}$ |