

CSE555: Introduction to Pattern Recognition

Midterm Exam

(100 points, Closed book/notes)

There are 5 questions in this exam.

Page 3 is the Appendix that contains some useful formulas.

1. (15pts) Bayes Decision Theory.

(a) (5pts) Assume there are c classes w_1, \dots, w_c , and one feature vector \mathbf{x} , give the Bayes rule for classification in terms of a priori probabilities of the classes and class-conditional probability densities of \mathbf{x} .

(b) (10pts) Suppose we have a two-classes problem ($A, \sim A$), with a single binary-valued feature ($\mathbf{x}, \sim \mathbf{x}$). Assume the prior probability $P(A) = 0.33$. Given the distribution of the samples as shown in the following table, use Bayes Rule to compute the values of posterior probabilities of classes.

	A	$\sim A$
\mathbf{x}	248	167
$\sim \mathbf{x}$	82	503


2. (25pts) Fisher Linear Discriminant.

(a) (5pts) What is the Fisher linear discriminant method?

(b) Given the 2-d data for two classes:

$\omega_1 = [(1, 1), (1, 2), (1, 4), (2, 1), (3, 1), (3, 3)]$ and

$\omega_2 = [(2, 2), (3, 2), (3, 4), (5, 1), (5, 4), (5, 5)]$ as shown in the figure:



- (10pts) Determine the optimal projection line in a single dimension.
- (10pts) Show the mapping of the points to the line as well as the Bayes discriminant assuming a suitable distribution.

3. (20pts) Suppose $p(x|w_1)$ and $p(x|w_2)$ are defined as follows:

$$p(x|w_1) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}, \quad \forall x$$

$$p(x|w_2) = \frac{1}{4}, \quad -2 < x < 2$$

- (a) (7pts) Find the minimum error classification rule $g(x)$ for this two-class problem, assuming $P(w_1) = P(w_2) = 0.5$.
- (b) (10pts) There is a prior probability of class 1, designated as π_1^* , so that if $P(w_1) > \pi_1^*$, the minimum error classification rule is to always decide w_1 regardless of x . Find π_1^* .
- (c) (3pts) There is no π_2^* so that if $P(w_2) > \pi_2^*$, we would always decide w_2 . Why not?
4. (20pts) Let samples be drawn by successive, independent selections of a state of nature w_i with unknown probability $P(w_i)$. Let $z_{ik} = 1$ if the state of nature for the k th sample is w_i and $z_{ik} = 0$ otherwise.

(a) (7pts) Show that

$$P(z_{i1}, \dots, z_{in} | P(w_i)) = \prod_{k=1}^n P(w_i)^{z_{ik}} (1 - P(w_i))^{1-z_{ik}}$$

(b) (10pts) Given the equation above, show that the maximum likelihood estimate for $P(w_i)$ is

$$\hat{P}(w_i) = \frac{1}{n} \sum_{k=1}^n z_{ik}$$

(c) (3pts) Interpret the meaning of your result in words.

5. (20pts) Consider an HMM with an explicit absorber state w_0 and unique null visible symbol v_0 with the following transition probabilities a_{ij} and symbol probabilities b_{jk} (where the matrix indexes begin at 0):

$$a_{ij} = \begin{pmatrix} 1 & 0 & 0 \\ 0.2 & 0.3 & 0.5 \\ 0.4 & 0.5 & 0.1 \end{pmatrix} \quad b_{jk} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0.7 & 0.3 \\ 0 & 0.4 & 0.6 \end{pmatrix}$$

- (a) (7pts) Give a graph representation of this Hidden Markov Model.
- (b) (10pts) Suppose the initial hidden state at $t = 0$ is w_1 . Starting from $t = 1$, what is the probability it generates the particular sequence $\mathbf{V}^3 = \{v_2, v_1, v_0\}$?
- (c) (3pts) Given the above sequence \mathbf{V}^3 , what is the most probable sequence of hidden states?

Appendix: Useful formulas.

- For a 2×2 matrix,

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

the matrix inverse is

$$A^{-1} = \frac{1}{|A|} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

- The *scatter matrices* \mathbf{S}_i are defined as

$$\mathbf{S}_i = \sum_{\mathbf{x} \in D_i} (\mathbf{x} - \mathbf{m}_i)(\mathbf{x} - \mathbf{m}_i)^t$$

where \mathbf{m}_i is the d -dimensional sample mean.

The *within-class scatter matrix* is defined as

$$\mathbf{S}_W = \mathbf{S}_1 + \mathbf{S}_2$$

The *between-class scatter matrix* is defined as

$$\mathbf{S}_B = (\mathbf{m}_1 - \mathbf{m}_2)(\mathbf{m}_1 - \mathbf{m}_2)^t$$

The solution for the \mathbf{w} that optimizes $J(\mathbf{w}) = \frac{\mathbf{w}^t \mathbf{S}_B \mathbf{w}}{\mathbf{w}^t \mathbf{S}_W \mathbf{w}}$ is

$$\mathbf{w} = \mathbf{S}_W^{-1}(\mathbf{m}_1 - \mathbf{m}_2)$$