


Querying Probabilistic Graphical Models

Sargur Srihari

srihari@cedar.buffalo.edu

Probabilistic Graphical Models



- Represent joint probability distributions over multiple variables
 - in terms of conditional probability distributions (CPDs)
 - Resulting in great savings in no of parameters needed

- CPDs: $P(X_i | pa(X_i))$

- Joint Distribution:

$$P(X) = P(X_1, \dots, X_n)$$

$$P(X) = \prod_{i=1}^N P(X_i | pa(X_i))$$

$$P(D, I, G, S, L) = P(D)P(I)P(G | D, I)P(S | I)P(L | G)$$

Queries of Interest

- Probabilistic Graphical Models represent joint probability distributions over multiple variables
- Their use is to answer queries of interest
- This is called “inference”
- What types of queries are there?

Query Types


1. Probability Queries
2. MAP Queries
 - Maximum a posteriori probability
 - Also called MPE (Most Probable Explanation)
3. Marginal MAP Queries

Probability Queries

- Most common type of query is a probability query
- Query has two parts
 - *Evidence*: a subset E of variables and their instantiation e
 - *Query Variables*: a subset Y of random variables in network
- **Inference Task: $P(Y|E=e)$**
 - *Posterior probability distribution* over values y of Y
 - *Conditioned* on the fact $E=e$
 - Can be viewed as Marginal over Y in distribution we obtain by conditioning on e
- **Marginal Probability Estimation**

$$P(Y = y_i | E = e) = \frac{P(Y = y_i, E = e)}{P(E = e)}$$

Example of Probability Query



$$P(Y = y_i | E = e) = \frac{P(Y = y_i, E = e)}{P(E = e)}$$

↑ Posterior Marginal ↑ Probability of Evidence

- Posterior Marginal Estimation: $P(I=i^1 | L=l^0, S=s^1) = ?$
- Probability of Evidence: $P(L=l^0, S=s^1) = ?$
 - Here we are asking for a specific probability rather than a full distribution

Computing the Probability of Evidence

Probability Distribution of Evidence

$$\begin{aligned}
 P(L, S) &= \sum_{D, I, G} P(D, I, G, L, S) && \text{Sum Rule of Probability} \\
 &= \sum_{D, I, G} P(D)P(I)P(G | D, I)P(L | G)P(S | I) && \text{From the Graphical Model}
 \end{aligned}$$

Probability of Evidence


$$P(L = l^0, s = s^1) = \sum_{D, I, G} P(D)P(I)P(G | D, I)P(L = l^0 | G)P(S = s^1 | I)$$

More Generally

$$P(E = e) = \sum_{X \setminus E} \prod_{i=1}^n P(X_i | pa(X_i)) | E = e$$

- An intractable problem
 - #P complete
- Tractable when tree-width is less than 25
 - Most real-world applications have higher tree-width
- Approximations are usually sufficient (hence sampling)
 - When $P(Y=y|E=e)=0.29292$, approximation yields 0.3

Probability of Random Correspondence (PRC)




- A type of query that can be asked of any distribution
- Probability that two independently chosen random samples X and Y from the same distribution will have the same value

$$\rho = P(Z = 1 | X, Y) = \sum_X \sum_Y P(Z = 1 | X, Y) P(X) P(Y)$$

- where $P(X) = P(Y)$ and $Z = 1$ if $X = Y + e$ and 0 otherwise
- If $e = 0$ then it is a case of exact matching where $\rho = \sum_X [P(X)]^2$

- **Conditional n PRC**

- Probability that evidence $X = e$ is found in database of size n
 $(1 - (1 - P(X = e))^n)$



MAP Queries


- Finding a high probability assignment to some subset of variables
- Most likely assignment to all non-evidence variables $W = \mathcal{X} - Y$

$$MAP(W | e) = \arg \max_w P(w, e)$$

Value of w for which $P(w, e)$ is maximum

- Difference from probability query
 - Instead of a probability we get the most likely value for all remaining variables

Example of MAP Query



$$MAP(A) = \arg \max_a P(A = a)$$

$$MAP(A, B) = \arg \max_{a, b} P(A, B)$$

$$= \arg \max_{a, b} P(A)P(B|A)$$

$$= \arg \max_{a, b} \{0.04, 0.36, 0.3, 0.3\}$$

$$= a^0, b^1$$


Note that individually most likely value a^1 is not in the most likely joint assignment

Marginal MAP Query

Diseases

a^0 a^1

0,4 0.6



Symptoms

A	b^0	b^1
a^0	0.1	0.9
a^1	0.5	0.5

We looked for highest joint probability assignment of disease and symptom

Can look for most likely assignment of disease variable only

Query is not all remaining variables but a subset of them

Y is query, evidence is $E=e$

Task is to find most likely assignment to Y :

$$MAP(Y|e) = \arg \max P(Y|e)$$

If $Z=X-Y-E$

$$MAP(Y | e) = \arg \max_Y \sum_Z P(Y, Z | e)$$

Contains both summations (like in probability queries)

And maximizations (like in MAP queries)

Marginal MAP Assignments

- They are not monotonic
- Most likely assignment $MAP(Y_1|e)$ might be completely different from assignment to Y_1 in $MAP(\{Y_1, Y_2\}|e)$
- Thus we cannot use a MAP query to give a correct answer to a marginal map query