

Markov Network Independencies


Sargur Srihari
srihari@cedar.buffalo.edu

Topics

- Markov Network Independencies
 - Basic Independencies
 - Independencies Revisited
 - From Distributions to Graphs

Markov network captures independencies of interactions

Misconception MN with factors



$$P(a,b,c,d) = \frac{1}{Z} \phi_1(a,b) \cdot \phi_2(b,c) \cdot \phi_3(c,d) \cdot \phi_4(d,a)$$

where

$$Z = \sum_{a,b,c,d} \phi_1(a,b) \cdot \phi_2(b,c) \cdot \phi_3(c,d) \cdot \phi_4(d,a)$$

Factorization of Distribution and Independences

P supports $(X \perp Y|Z)$ iff we can write distribution as $P(X) = \Phi_1(X,Z) \Phi_2(Y,Z)$

$$P(A,B,C,D) = \left[\frac{1}{Z} \phi_1(A,B) \cdot \phi_2(B,C) \right] \phi_3(C,D) \cdot \phi_4(D,A)$$

Factor with $\{B, \{A, C\}\}$


Factor with $\{D, \{A, C\}\}$

Therefore $(B \perp D|A, C)$

Similarly $(A \perp C|B, D)$

Basic Independencies

- As in Bayesian Networks, graph structure in a Markov network encodes a set of independence assumptions
- In a MN Probabilistic influence flows along the undirected paths in the graph and “blocked” if we condition on intervening nodes



Active Paths, Separation, Global Independence

H is a Markov network

X_1, \dots, X_k is a path in H

Z : set of observed variables

1. Path is active given Z if none of the X_i is in Z


- If B and D are observed, path A, C is inactive

2. Set of nodes Z separates sets X and Y

- If no active path between any node $X \in X$ and $Y \in Y$

3. Global independencies associated with H are

- $I(H) = \{(X \perp Y | Z) : sep_H(X, Y | Z)\}$
- Independencies in $I(H)$ guaranteed to hold for every distribution P over H



$$I(H) = \{(A \perp C | B, D), (B \perp D | A, C)\}$$

Formalizing independencies in MNs and distributions

- Gibbs Distribution

- A distribution P_Φ is a Gibbs distribution parameterized by a set of factors $\Phi = \{\phi_1(D_1), \dots, \phi_K(D_K)\}$
- If defined as follows

$$P_\Phi(X_1, \dots, X_n) = \frac{1}{Z} \tilde{P}(X_1, \dots, X_n)$$

where

$$\tilde{P}(X_1, \dots, X_n) = \prod_{i=1}^m \phi_i(D_i)$$

D_i are sets of random variables

is an unnormalized measure and

$$Z = \sum_{X_1, \dots, X_n} \tilde{P}(X_1, \dots, X_n)$$

Soundness of Separation

- Theorem 1:
 - Let P be a distribution over $\chi = \{X_1, \dots, X_n\}$ and H is a Markov structure over χ
 - If P is a Gibbs distribution that factorizes over H , then H is an I-map for P
 - Factorize means every D_i in H is a clique
- Theorem 2: Hammersley-Clifford (other direction)
 - if H is an I-map for P then P factorizes over H
 - Holds only for positive distributions ($P > 0$ for every assignment)

Positive Distribution


- A distribution P is said to be positive if for all events $\alpha \in S$ such that $\alpha \neq \Phi$ we have that $P(\alpha) > 0$
- A non-positive distribution
- 16 possible values
 - Distribution P : 8 have value $1/8$ rest are zero

X_1	X_2	X_3	X_4	$P(X)$
0	0	0	0	1/8
1	0	0	0	1/8
1	1	0	0	1/8
1	1	1	0	1/8
0	0	0	1	1/8
0	0	1	1	1/8
0	1	1	1	1/8
1	1	1	1	1/8

Rest 8 probs are 0

Non-positive distribution consistent with H

- Four binary random variables
- Independencies in graph H:
- Consider global independencies
- E.g. $X_1 \perp\!\!\!\perp X_3 \mid X_2, X_4$
 - implies $(X_1 \perp\!\!\!\perp X_3 \mid X_2, X_4)$
- P also satisfies this
 - For the assignment $X_2=1, X_4=0$
 - $P(X_1=1 \mid X_2=1, X_4=0)=1$
 - Rest are zero
 - Thus X_1 is independent of X_3




X_1	X_2	X_3	X_4	$P(\mathbf{X})$
0	0	0	0	1/8
1	0	0	0	1/8
1	1	0	0	1/8
1	1	1	0	1/8
0	0	0	1	1/8
0	0	1	1	1/8
0	1	1	1	1/8
1	1	1	1	1/8

Rest 8 probs are 0

- Global independencies hold \rightarrow H is an I-map for P
- But P does not factorize according to H (Proof by contradiction)

Graphical Model as Filter



$p(x)$ is allowed to pass through only if
It satisfies independencies in graph
This set is denoted DF or UF (for BN or MN)

- UI is set of distributions that are consistent with set of conditional independence statements read from the graph using graph separation
- UF are set of distributions that can be expressed as factorization of the form

$$p(\mathbf{x}) = \frac{1}{Z} \prod_c \psi_c(\mathbf{x}_c)$$

- *Hammersley-Clifford theorem* states that UI and UF are identical

Independencies in Bayesian Networks

- Bayesian networks have two types of independencies
 - Local independencies
 - Each node is independent of its non-descendants given its parents
 - Global independencies
 - Induced by d-separation
- These two sets of independencies are equivalent

Three Independencies of Markov Networks

- Two local independencies
 - Pairwise $I_p(H)$
 - Markov Blanket $I_l(H)$
- Global independency $I(H)$
 - Identify three sets of nodes A , B and C
 - To test conditional independence property
$$A \perp B | C$$
 - Consider all possible paths from nodes in set A to nodes in set B
 - If all such paths pass through one or more nodes in C then path is blocked and independence holds


Pairwise Independency $I_p(H)$

- Weakest independency
 - Whenever two variables are directly connected
 - Potential of being correlated not mediated by other variables
 - When two variables are not directly linked
 - There must be a way of rendering them conditionally independent
- If H is a Markov network its *pairwise independencies* are

$$I_p(H) = \{(X \perp Y | \chi - \{X, Y\}) : X - Y \notin H\}$$

- In the example


$$I_p(H) = \{(A \perp C | B, D), (B \perp D | A, C)\}$$



When X, Y are not directly connected they are independent given other variables

Markov Blanket Independency $I_I(H)$

- Analogous to local independencies in Bayesian networks
 - We can block all influences by conditioning on its immediate neighbors
 - Node is conditionally independent of all nodes given its immediate neighbors
- For graph H the Markov blanket of X in H is the set of neighbors of X in H
- Local independencies associated are
 - $I_I(H) = \{(X \perp \chi - \{X\} - MB_H(X) \mid MB_H(X)) : X \in \chi\}$



Relationship between Markov properties

- Three independencies of network structure H
- $I_p(H)$ is strictly weaker than $I_l(H)$ is strictly weaker than $I(H)$
- For positive distributions all three are equivalent

Separation in Markov Networks

- Markov network encodes a set of conditional independencies

- Probabilistic influence flows

- in undirected paths

- Blocked if we condition on intervening nodes

- Every path from any node in A to B passes through C

- No explaining away


- Testing for independence simpler than in directed graphs

- Alternative view


- Remove all nodes in set C together with all their connecting links

- If no paths from A to B then conditional independence holds

- Markov blanket




- A node is conditionally independent of all other nodes conditioned only on its neighbors



Factorization Properties

- Factorization rule corresponds to conditional independence test
- Notion of locality needed
- Consider two nodes x_i and x_j not connected by a link
 - They are conditionally independent given all other nodes in graph
 - Because there is no direct path between them and
 - All other paths pass through nodes that are observed and hence those paths are blocked
 - Expressed as

$$p(x_i, x_j \mid \mathbf{X} \setminus \{i, j\}) = p(x_i \mid \mathbf{X} \setminus \{i, j\}) p(x_j \mid \mathbf{X} \setminus \{i, j\})$$
 - Where $\mathbf{X} \setminus \{i, j\}$ denotes set \mathbf{x} of all variables with x_i and x_j removed
- For conditional independence to hold
 - factorization is such that x_i and x_j do not appear in the same factor
 - No path between them other than going through others
 - leads to graph concept of clique



From Distributions to Graphs

- How to encode independencies in a given distribution P using a graph structure
 - A complete graph (all nodes connected) encodes no independencies
 - How can we construct a minimal I-map
- Two equivalent approaches
 - Based on pairwise Markov independencies
 - Based on local independencies

P to H using Pairwise Independencies

- We must add edges between all pairs of nodes X, Y such that
 - P does not support $(X \perp Y | \chi - \{X, Y\})$
 - i.e., If two variables are not independent given other variables, then add an edge

P to H using Local Independencies

- Use notion of Markov Blanket defined for P
- For each variable X we define neighbors of Z to be a minimal set of nodes Y that render X independent of rest of nodes
- U is a Markov Blanket of X in distribution P if $X \perp_{\mathcal{X}-\{X\}-U|U} \in I(P)$
- Define graph H by introducing edge $\{X, Y\}$ for all $X, Y \in MB_P(X)$

