

# Forward Sampling

Sargur Srihari  
srihari@buffalo.edu

# Topics

- Forward Sampling
- Sampling from a Bayesian Network
- Sampling from a Discrete Distribution
- Use of samples
- Analysis of Error
- Conditional Probability Queries

# Forward Sampling: Plan of Discussion

- Forward sampling

- It is the simplest particle generation approach
- We generate samples  $\xi[1], \dots, \xi[M]$  from  $P(\boldsymbol{\chi})$

$\xi$  denotes a full assignment to variables in  $\boldsymbol{\chi}$ , i.e.,  $\boldsymbol{\chi} \in \text{Val}(\boldsymbol{\chi})$


- Plan of discussion

- Generating particles from  $P_{\mathcal{B}}(\boldsymbol{\chi})$  by sampling a BN
- No of particles needed to get a good approximation of the expectation of a target function  $f$
- Difficulties in generating samples from posterior  $P_{\mathcal{B}}(\boldsymbol{\chi}|e)$ 
  - In undirected models even generating a sample from the prior distribution is a difficult task

# Sampling from a Bayesian Network

- A very simple process
- Sample nodes in some order so that by the time we sample a node we have values for all of its parents
- We can then sample from the distribution specified by the CPD
- Need ability to sample from the distributions underlying the CPD
  - Straightforward for discrete case
  - Subtler for continuous measures


# Ex: Forward (Ancestral) Sampling



1 Sample  $D \{0.6 \text{ vs } 0.4\} \rightarrow d^0$   
 2 Sample  $I \{0.7 \text{ vs } 0.3\} \rightarrow i^0$   
 3 Sample  $G \{0.3, 0.4, 0.3\} \rightarrow g^2$   
 etc, etc

# Sampling from a Discrete Distribution

- Split  $[0,1]$  interval into bins whose sizes are determined by the probabilities  $P(x^i)$ ,  $i=1,\dots,k$ 
  - Partition the interval into  $k$  subintervals
- Generate a sample  $s$  uniformly from the interval
- If  $s$  is in the  $i^{\text{th}}$  interval then sampled value is  $x^i$
- Example: sample from  $P(x=\{1,2,3,4\})=\{0.3,0.1,0.4,0.2\}$



# Using samples to find expectations

- To compute an expectation  $E[f(\mathbf{x})] = \sum_{\mathbf{x}} p(\mathbf{x})f(\mathbf{x})$ 
  - Using basic convergence bounds, we know that from a set of particles  $\mathcal{D} = \{\xi[1], \dots, (\xi[M])\}$  generated via sampling we can estimate it as

$$\hat{E}_D(f) = \frac{1}{M} \sum_{m=1}^M f(\xi[m])$$

- In the case where our task is to compute  $P(\mathbf{y})$ 
  - for event  $Y=\mathbf{y}$  relative to  $P$  for some  $Y \subseteq X$  and  $\mathbf{y} \in \text{Val}(Y)$ 
    - this estimate is simply the fraction of particles where we have seen the event  $\mathbf{y}$

$$\hat{P}_D(\mathbf{y}) = \frac{1}{M} \sum_{m=1}^M I\{\mathbf{y}[m] = \mathbf{y}\}$$

$\mathbf{y}[m]$  denotes  $\xi[m] \mathbf{Y}$   
i.e., assignment to  $\mathbf{Y}$  in the particle  $\xi[m]$

# Analysis of Error

- Quality of the estimate depends heavily on the no. of particles generated
- How many particles are needed to obtain a certain performance guarantee?
- Focus analysis on the case where we need to estimate  $P(\mathbf{y})$

# How many samples are required?

## 1. From Hoeffding bound

- Error is bounded by  $\varepsilon$  with probability of at least  $1-\delta$
- The estimator with  $(\varepsilon, \delta)$  reliability is

$$\begin{aligned} &\text{For } M \text{ ind Bernoulli trials with prob (success)}=p \\ &S_D = X[1] + \dots + X[M], \quad T_D = (1/M)S_D \\ &P(T_D > p + \varepsilon) \leq \exp(-2M\varepsilon^2) \\ &P(T_D < p - \varepsilon) \leq \exp(-2M\varepsilon^2) \end{aligned}$$

$$M \geq \frac{\ln(2 / \delta)}{2\varepsilon^2}$$

## 2. From Chernoff bound

- For a given  $\varepsilon$ , no. of samples needed to guarantee a certain error probability  $\delta$  is

$$M \geq 3 \frac{\ln(2 / \delta)}{P(y)\varepsilon^2}$$

- $M$  grows inversely with probability  $P(y)$
- Thus, cannot determine no. of samples needed

# Conditional Probability Queries

- So far we have discussed the problem of estimating marginal probabilities
  - i.e., Probability of event  $Y=y$  relative to the original joint distribution
- In general we are interested in conditional distributions of the form  $p(\mathbf{y}|\mathbf{E}=\mathbf{e})$
- This estimation task is significantly harder
- One approach is called Rejection sampling

# Rejection Sampling

- Generate samples from posterior  $P(\boldsymbol{\chi}|\boldsymbol{e})$ 
  - Can do this by generating samples  $\boldsymbol{x}$  from  $P(\mathbf{X})$
  - Reject any sample that is not compatible with  $\boldsymbol{e}$
  - Resulting samples are from posterior  $P(\boldsymbol{\chi}|\boldsymbol{e})$
- Problem: no of unrejected particles is small
  - Expected no of unrejected from original  $M$  is  $MP(\boldsymbol{e})$ 
    - Ex: if  $P(\boldsymbol{e})=0.001$  &  $M=10,000$ , only 10 unrejected samples
  - Conversely, to obtain at least  $M^*$  unrejected particles, we need to generate on an average  $M=M^*/P(\boldsymbol{e})$  samples from  $P(\mathbf{X})$

# Low probability evidence is common

- In many applications, low probability evidence is the rule rather than the exception
- Example of medical diagnosis
  - Any set of symptoms typically has a low probability
    - As no of variables  $k$  increases, probability of evidence  $k$  decreases exponentially with  $k$

## Indirect Approach also has limitation

- Alternative approach to determining  $P(\mathbf{y}|\mathbf{E}=\mathbf{e})$  is to use a separate estimator for  $P(\mathbf{e})$  and for  $P(\mathbf{y},\mathbf{e})$  and then compute the ratio
  - If both quantities gave low relative error then their ratio also will have low relative error
- No. of samples required to get a low relative error also grows linearly with  $1/P(\mathbf{e})$