

Probability of Random Correspondence for Fingerprints

Chang Su and Sargur N. Srihari

Computer Science and Engineering Department
University at Buffalo, Buffalo NY 14228, USA
{changsu, srihari}@cedar.buffalo.edu,

Abstract. Individuality of fingerprints can be quantified by computing the probabilistic metrics for measuring the degree of fingerprint individuality. In this paper, we present a novel individuality evaluation approach to estimate the probability of random correspondence (PRC). Three generative models are developed respectively to represent the distribution of fingerprint features: ridge flow, minutiae and minutiae together with ridge points. A mathematical model that computes the PRCs are derived based on the generative models. Three metrics are discussed in this paper: (i) PRC of two samples, (ii) PRC among a random set of n samples (n PRC) and (iii) PRC between a specific sample among n others (specific n PRC). Experimental results show that the theoretical estimates of fingerprint individuality using our model consistently follow the empirical values based on the NIST4 database.

1 Introduction

We consider the evaluation of the relevant probabilities in a commonly used forensic modality – that of fingerprints. Their use in human identification has been based on two premises, that, (i) they do not change with time and (ii) they are unique for each individual. While in the past, identification based on fingerprints had been accepted by courts, more recently their use has been questioned under the basis that the premises stated above have not been objectively tested and error rates not been scientifically established [1]. Though the first premise has been accepted, the second one on individuality is being challenged.

Fingerprint individuality studies date back to the late 1800s. More than twenty models have been proposed to establish the improbability of two random people (or fingers) have the same fingerprint [2]. These models can be classified into five different categories: grid-based [3, 4], ridge-based [4], fixed probability [5], relative measurement [6] and generative [7–9]. All models try to quantify the uniqueness property, e.g., the probability of false correspondence.

Features for representing fingerprints are classified into three types [10]. Level 1 features provide class-characterization of fingerprints based on ridge flow. Fig.1 shows five primary classes: whorl, left loop, right loop, arch and tent. Level 2 features, which are more useful for identification, are also known as minutiae. The minutiae correspond to ridge endings and ridge bifurcations. A minutia is




Fig. 1. Examples of five main types of ridge flow, referred to as Level 1 features: (a) arch, (b) left loop, (c) right loop, (d) tented arch, and (e) whorl.

represented by its location and direction; direction is determined by the ridge ending at the location (Figure 2(a)). Automatic fingerprint matching algorithms use minutiae as the salient features [11], since they are stable and are reliably extracted. For some cases in which minutiae can not provide sufficient information, ridge points are also taken into account. The ridge points are shown in Figure 2(b). Level 3 features, such as pores and scars are ancillary features. In this paper, only level 1 and level 2 features are considered. Our goal is to model the distribution of fingerprint features and then estimate the individuality of fingerprints based on the models.




Fig. 2. Representation of fingerprints using minutiae and ridgepoints: (a) locations of ridge endings and ridge bifurcations are indicated by circles, and (b) ridge points are indicated by stars in a skeletonized fingerprint image.

This paper is the subsequence of our work on generative models on fingerprints [9]. We measure individuality by 3 metrics: PRC, n PRC and specific PRC on level 1 and level 2 features. A novel PRC calculate algorithm is proposed to get rid of the dependence on the fingerprint matchers. The rest of this paper is organized as follows: Section 2 describes the generative models proposed for the ridge flow types, minutiae only and a combination- representative ridges. Methods for estimating individuality and experimental results are described in Section 3. Summary is given in Section 4.

2 Generative Models for Fingerprints

2.1 Distribution of Ridge Flow Types

A simple distribution of the Level 1 ridge flow types is obtained by counting the relative frequency of each of the primary and secondary types in a fingerprint database. In one such evaluation [16] loops account for 64% of the fingers, with the secondary types being: 30% left loops, 27% right loops and 7% double loops. Arches account for 18% of the primary types, with the secondary types being: plain arches (13%) and tented arches (5%). Whorls account for the remainder of the Level 1 types (19%).

2.2 Distribution of Minutiae

Each minutia is represented as $\mathbf{x} = (s, \theta)$ where $s = (x_1, x_2)$ is its location and θ its direction. The distribution of minutiae location is shown in Fig. 3(a). The minutiae is taken from 2000 fingerprints in NIST4 database.




Fig. 3. Model for minutiae distribution: (a) Distribution of minutia location from NIST special database, (b) Three-dimensional plot of mixture Gaussian model for minutia location with three components, (c) von Mises distributions of minutiae orientation for each of the three components, where the green curve corresponds to the upper cluster, blue the lower left cluster and red the lower right cluster.

Since minutia location has a multimodal distribution as shown in Fig.3(a), a mixture of K Gaussians is a natural approach. Furthermore since minutiae orientation is a periodic variable, it is modeled by a *circular normal* or von Mises distribution [12, 13]. Such a model is better than mixtures of hyper-geometric and binomial distributions [7, 8]. We write the distribution of minutiae as

$$p(\mathbf{x}|\Theta) = \sum_{k=1}^K \pi_k \cdot \mathcal{N}(s|\mu_k, \Sigma_k) \cdot \mathcal{V}(\theta|\nu_k, \kappa_k), \quad (1)$$

where K is the number of mixture components, π_k are non-negative component weights that sum to one, $\mathcal{N}(s|\mu_k, \Sigma_k)$ is the bivariate Gaussian probability density function of minutia with mean μ_k and covariance matrix Σ_k , $\mathcal{V}(\theta|\nu_k, \kappa_k)$ is the von Mises probability density function of minutiae orientation with mean angle ν_k and precision κ_k , and $\Theta = \{\pi_k, \mu_k, \Sigma_k, \nu_k, \kappa_k, \rho_k\}$ where $k = 1, 2, \dots, K$ is the set of all parameters of the k Gaussian and von Mises distributions.

Rather than using the standard form of the von Mises distribution for the range $[0, 2\pi]$, since minutiae orientations are represented as being in the range $[0, \pi)$, we use the alternate form [13] as follows

$$\mathcal{V}(\theta|\nu_k, \kappa_k, \rho_k) = \rho_k v(\theta) \cdot I\{0 \leq \theta < \pi\} + (1 - \rho_k) v(\theta - \pi) \cdot I\{\pi \leq \theta < 2\pi\} \quad (2)$$

where $I\{A\}$ is the indicator function of the condition A ,

$$v(\theta) \equiv v(\theta|\nu_k, \kappa_k) = \frac{1}{\pi I_0(\kappa_k)} \exp[\kappa_k \cos 2(\theta - \nu_k)], \quad (3)$$

minutiae arising from the k^{th} component have directions that are either θ or $\theta + \pi$ and the probabilities associated with these two occurrences are ρ_k and $1 - \rho_k$ respectively.

Since fingerprint ridges flow smoothly with very slow direction changes, direction of neighboring minutiae are strongly correlated, i.e., minutiae that are spatially close tend to have similar directions with each other. However, minutiae in different regions of a fingerprint tend to be associated with different region-specific minutiae directions thereby demonstrating independence [14, 15]. The model allows ridge orientations to be different at different regions (different regions can be denoted by different components) while it makes sure that nearby minutiae have similar orientations (as nearby minutiae will belong to the same component). The parameters for generative models are learned from NIST4 fingerprint database. We show the results in Figure 3(b) 3(c).

2.3 Distribution of Representative Ridges

In the model just discussed, only minutiae was used in the framework of generative models for fingerprints. Models based purely on minutiae may be sufficient to model biometric scenarios where finger-prints are obtained in controlled conditions [7][8], but are insufficient to model forensic scenarios where latent prints are lifted off of surfaces. In such cases, ridge details provide vital information about fingerprints. Verification systems using minutiae together with ridge information are more accurate than using minutiae alone. Also, any generative model that makes use of ridge details can only be a better representation of the generative model for fingerprints. With this motivation, the distribution of ridge information is embedded into generative models [9].

Since nearby ridges always maintain similar trend, too much redundancy would be involved if all the ridges are modeled. We only consider ridges where the minutiae lie. These ridges are called *representative ridges*. A representative ridge contains two parts, the minutiae and the following ridge. The latter part

is represented as a set of ridge points sampled at equal interval of inter ridge width [18]. The length of the ridge l_r is defined as the number of ridge points that could be sampled on the ridge. Three types of ridges are defined: (i) short: $l_r \leq L/3$, (ii) medium: $L/3 < l_r \leq 2L/3$ and (iii) long: $l_r > 2L/3$, where L is the average ridge length of the top 10% longest representative ridges in the fingerprints database, e.g. in NIST4, L is 18. By choosing the value of maximum ridge length L as the average of long ridges in the database, unusually long ridges caused by artifacts is avoided. The three possible ridge length types can be associated with any representative ridge. Without loss of generality, we can assume that there exist only three possible ridge length types corresponding to a representative ridge. For ridges with different lengths, different ridge points are picked as anchors. For medium ridges, $\lfloor L/3 \rfloor^{th}$ ridge point is picked and for long ridges, both $\lfloor L/3 \rfloor^{th}$ and $\lfloor 2L/3 \rfloor^{th}$ are picked. No ridge point is chosen for short ridges. The rationale for choosing these two ridge points is described in [18].

Let $\mathbf{x} = \{\mathbf{x}_m, \mathbf{x}_r\}$ denote the feature vector of a representative ridge, where minutiae feature vector \mathbf{x}_m is given by $\{s_m, \theta_m\}$ and ridge feature vector \mathbf{x}_r is represented by a ridge point set given by $\{\mathbf{x}_{ri} : i \in A \wedge i \leq l_r\}$ where \mathbf{x}_{ri} is the feature vector of the i^{th} ridge point, A is the anchor point index set $\{\lfloor L/3 \rfloor, \lfloor 2L/3 \rfloor\}$ and l_r is the length of the ridge. In contrast to the model for minutiae, ridge points are represented in polar coordinates as shown in Figure 4. The location of ridge point s_{ri} is given by $\{r_i, \phi_i\}$ and the direction of ridge point is θ_i . Thus the ridge point \mathbf{x}_{ri} is represented as the combination of location and direction $\{r_i, \phi_i, \theta_i\}$ where r_i is the distance from the i^{th} ridge point to the minutia, ϕ_i is the positive angle required to reach the i^{th} ridge point from the polar axis.




Fig. 4. Representation of ridge points in polar coordinates.

The generative model is based on the distribution of representative ridges. Mixture distributions consisting of K_i components, $i = 1, 2, 3$, is used to model representative ridges of three ridge length types. Each component is distributed according to the density of the minutiae and density of ridge points. Assuming that

the minutiae and ridge points are independent, representative ridge distribution is given by

$$p(\mathbf{x}|\Theta) = \begin{cases} p(l_r) \cdot \sum_{g=1}^{K_1} \pi_k p_k(s_m, \theta_m | \Theta_k) & l_r \leq \frac{L}{3} \\ p(l_r) \cdot \sum_{k=1}^{K_2} \pi_k p_k(s_m, \theta_m | \Theta_k) \cdot p_k(r_{\lfloor \frac{L}{3} \rfloor}, \phi_{\lfloor \frac{L}{3} \rfloor}, \theta_{\lfloor \frac{L}{3} \rfloor} | \Theta_k) & \frac{L}{3} < l_r < \frac{2L}{3} \\ p(l_r) \cdot \sum_{k=1}^{K_3} \pi_k p_k(s_m, \theta_m | \Theta_k) \cdot p_k(r_{\lfloor \frac{L}{3} \rfloor}, \phi_{\lfloor \frac{L}{3} \rfloor}, \theta_{\lfloor \frac{L}{3} \rfloor} | \Theta_k) \cdot p_k(r_{\lfloor \frac{2L}{3} \rfloor}, \phi_{\lfloor \frac{2L}{3} \rfloor}, \theta_{\lfloor \frac{2L}{3} \rfloor} | \Theta_k) & l_r \geq \frac{2L}{3} \end{cases} \quad (4)$$

The first condition corresponds to minutiae alone, the second to minutia and one ridge point, and the third to minutia and two ridge points. $p_k(s_m, \theta_m | \Theta_k)$ is the component distribution for minutiae location s_m and direction θ_m and $p_k(r_i, \phi_i, \theta_i | \Theta_k)$ is the component distribution for the i^{th} ridge point. They are defined as in (5) and (6) respectively.

$$p_k(s_m, \theta_m | \Theta_k) = \mathcal{N}(s_m | \mu_{mk}, \Sigma_{mk}) \cdot \mathcal{V}(\theta_m | \nu_{mk}, \kappa_{mk}, \rho_{mk}) \quad (5)$$

$$p_k(r_i, \phi_i, \theta_i | \Theta_k) = p_k(r_i, \phi_i | \mu_{ik}, \Sigma_{ik}, \nu_{ik}^\phi, \kappa_{ik}^\phi, \rho_{ik}^\phi) \cdot \mathcal{V}(\theta_i | \nu_{ik}^\theta, \kappa_{ik}^\theta, \rho_{ik}^\theta) \quad (6)$$

where $p_k(s_m, \theta_m | \Theta_k)$ can be calculated by Eq. 1 and $p_k(r_i, \phi_i, \theta_i | \Theta_k)$ is the product of the probabilities of ridge point locations and directions, where $\mathcal{V}(\theta_i | \nu_{ik}^\theta, \kappa_{ik}^\theta, \rho_{ik}^\theta)$ presents the distribution of the ridge point direction, θ_i is the direction of the i^{th} ridge point and $p_k(r_i, \phi_i | \mu_{ik}, \Sigma_{ik}, \nu_{ik}^\phi, \kappa_{ik}^\phi, \rho_{ik}^\phi)$ is the distribution of ridge point location given by

$$p_k(r_i, \phi_i | \mu_{ik}, \Sigma_{ik}, \nu_{ik}^\phi, \kappa_{ik}^\phi, \rho_{ik}^\phi) = \mathcal{N}(r_i | \mu_{ik}, \sigma_{ik}) \cdot \mathcal{V}(\phi_i | \nu_{ik}^\phi, \kappa_{ik}^\phi, \rho_{ik}^\phi) \quad (7)$$

where $\mathcal{N}(r_i | \mu_{ik}, \sigma_{ik})$ is a univariate Gaussian distribution whose mean μ_{ik} and variance σ_{ik} are learnt from a fingerprint database.

3 Evaluation of PRCs

As measures of individuality, three probabilities can be defined: PRC, n PRC and Specific n PRC. These definitions are further described below.

1. PRC: probability that two randomly chosen samples have the same measured value \mathbf{x} within specified tolerance ϵ .
2. n PRC: the probability that among a set of n samples, some pair have the same value \mathbf{x} , within specified tolerance, where $n \geq 2$. Since there are $\binom{n}{2}$ pairs involved this probability is higher than PRC. Note that when $n = 2$, PRC= n PRC.
3. Specific n PRC: the probability that in a set of n samples, a specific one with value \mathbf{x} coincides with another sample, within specified tolerance. Since we are trying to match a specific value \mathbf{x} , this probability depends on the probability of \mathbf{x} and is generally smaller than PRC. The exact relationship with respect to PRC depends on the distribution of \mathbf{x} .

We note that the first two measures, PRC and n PRC characterize the forensic modality as described by a set of measurements and furthermore, the second is a function of the first. The third measure specific n PRC characterizes specific evidence, e.g., a specimen. It is dependent on the specific value as well as the distribution of the measurement.

3.1 Ridge Flow Types

Assuming that fingerprints are distinguished by 6 secondary types, the PRC for ridge flow types is calculated by definition of PRC. Given type frequencies mentioned in section 2.1, we have PRC value $p_\epsilon = 0.2233$. The n PRC and specific n PRC can be simply computed from PRC value. Table 1 and table 2 show the n PRC and Specific n PRC with different n .

Table 1. Ridge Flow Types: n PRC

| n | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|---------|--------|--------|--------|--------|--------|---|---|
| n PRC | 0.2233 | 0.5314 | 0.7805 | 0.9201 | 0.9774 | 1 | 1 |

Table 2. Ridge Flow Types: Specific n PRC belong to class LL = left loop, RL = right loop, DL = double loop, PA = plain arch, TA = tented arch and W = whorl.

| n | Specific n PRC | | | | | |
|-----|------------------|--------|--------|--------|--------|--------|
| | LL | RL | DL | PA | TA | W |
| 10 | 0.9596 | 0.9411 | 0.4796 | 0.7145 | 0.3698 | 0.8499 |
| 20 | 0.9989 | 0.9975 | 0.7481 | 0.9291 | 0.6226 | 0.9818 |

Level 1 features are clearly broad class characteristics which are useful for exclusion of individual fingers but not by themselves useful for the tasks of verification, identification and individualization.

3.2 Minutiae Only

To compute the PRCs for minutiae, we first define correspondence, or match, between two minutiae. Let $\mathbf{x}_a = (s_a, \theta_a)$ and $\mathbf{x}_b = (s_b, \theta_b)$ be a pair of minutiae. The minutiae are said to correspond if for tolerance $\epsilon = [\epsilon_s, \epsilon_\theta]$,

$$|s_a - s_b| \leq \epsilon_s \text{ and } |\theta_a - \theta_b| \leq \epsilon_\theta \quad (8)$$

where $|s_a - s_b|$ is the Euclidean distance between the minutiae location $s_a = (x_{a1}, x_{a2})$ and $s_b = (x_{b1}, x_{b2})$.

Then, the probability that a random minutia \mathbf{x}_a would match a random minutia \mathbf{x}_b is given by

$$\begin{aligned} p_\epsilon(\mathbf{x}) &= p(|\mathbf{x}_a - \mathbf{x}_b| \leq \epsilon | \Theta) \\ &= \int_{\mathbf{x}_a} \int_{|\mathbf{x}_a - \mathbf{x}_b| \leq \epsilon} p(\mathbf{x}_a | \Theta) p(\mathbf{x}_b | \Theta) d\mathbf{x}_a d\mathbf{x}_b \end{aligned} \quad (9)$$

where Θ is the set of parameters describing the distribution of the minutiae location and direction.

Finally, the PRC, or the probability of matching at least \hat{m} pairs of minutiae within ϵ between two randomly chosen fingerprint f_1 and f_2 is calculated as

$$p_\epsilon(\hat{m}, m_1, m_2) = \binom{m_1}{\hat{m}} \binom{m_2}{\hat{m}} \hat{m}! \cdot p_\epsilon(\mathbf{x})^{\hat{m}} (1 - p_\epsilon(\mathbf{x}))^{(m_1 - \hat{m}) \cdot (m_2 - \hat{m})} \quad (10)$$

where m_1 and m_2 are numbers of minutiae in fingerprints f_1 and f_2 , $p_\epsilon(\mathbf{x})^{\hat{m}}$ is the probability of matching \hat{m} specific pairs of minutiae between f_1 and f_2 , $(1 - p_\epsilon(\mathbf{x}))^{(m_1 - \hat{m}) \cdot (m_2 - \hat{m})}$ is the probability that none of minutiae pair would match between the rest of minutiae in f_1 and f_2 and $\binom{m_1}{\hat{m}} \binom{m_2}{\hat{m}} \hat{m}!$ is the number of different match sets that can be paired up.

Given n fingerprints and assuming that the number of minutiae in a fingerprint m can be modeled by the distribution $p(m)$, the general PRCs $p(n)$ is given by

$$p(n) = 1 - \bar{p}(n) = 1 - (1 - p_\epsilon)^{\frac{n(n-1)}{2}} \quad (11)$$

where p_ϵ is the probability of matching two random fingerprint from n fingerprints. If we set the tolerance in terms of number of matching minutiae to \hat{m} , p_ϵ is calculated by

$$p_\epsilon = \sum_{m'_1 \in M_1} \sum_{m'_2 \in M_2} p(m'_1) p(m'_2) p_\epsilon(\hat{m}, m'_1, m'_2) \quad (12)$$

where M_1 and M_2 contain all possible numbers of minutiae in one fingerprint among n fingerprints, and $p_\epsilon(\hat{m}, m'_1, m'_2)$ can be calculated by Eq. 10.

Given a specific fingerprint f , the specific n PRCs can be computed by

$$p(f, n) = 1 - (1 - p(f))^{n-1} \quad (13)$$

where $p(f)$ is the probability that \hat{m} pairs of minutiae are matched between the given fingerprint f and a randomly chosen fingerprint from n fingerprints.

$$\begin{aligned} p(f) &= \sum_{m' \in M} p(m') \binom{m'}{\hat{m}} \cdot \sum_{i=1}^{\binom{m_f}{\hat{m}}} p(f_i) \\ &= \sum_{m' \in M} p(m') \binom{m'}{\hat{m}} \cdot \sum_{i=1}^{\binom{m_f}{\hat{m}}} \prod_{j=1}^{\hat{m}} p(\mathbf{x}_{ij} | \Theta) \end{aligned} \quad (14)$$

where M contains all possible numbers of minutiae in one fingerprint among n fingerprints, $p(m')$ is the probability of a fingerprint having m' minutiae in n fingerprints, m_f is the number of minutiae in the given fingerprint f , minutiae set $f_i = (\mathbf{x}_{i1}, \mathbf{x}_{i2}, \dots, \mathbf{x}_{im})$ is the subset of the minutiae set of given fingerprint and $p(f_i)$ is the joint probability of minutiae set f_i based on learned generative model.

3.3 Minutiae with Ridge Points

When ridge information is considered, a representative ridge is denoted by $\mathbf{x} = \{\mathbf{x}_m, \mathbf{x}_r\}$, where $\mathbf{x}_r = \{\mathbf{x}_r^i : i \in \{[L/3], [2L/3]\} \wedge i \leq l_i\}$. The representative ridge \mathbf{x}_a matches the representative ridge \mathbf{x}_b with tolerance ϵ if

$$|\mathbf{x}_{ma} - \mathbf{x}_{mb}| \leq \epsilon_m \wedge |\mathbf{x}_{ra} - \mathbf{x}_{rb}| \leq \epsilon_r \quad (15)$$

where $|\mathbf{x}_{ma} - \mathbf{x}_{mb}| \leq \epsilon_m$ is define by Eq.8 and $|\mathbf{x}_{ra} - \mathbf{x}_{rb}| \leq \epsilon_r$ is defined as

$$|\mathbf{x}_{ra} - \mathbf{x}_{rb}| \leq \epsilon_r \equiv (\forall i \in A) |r_a^i - r_b^i| \leq \epsilon_r \wedge |\phi_a^i - \phi_b^i| \leq \epsilon_\phi \wedge |\theta_a^i - \theta_b^i| \leq \epsilon_\theta \quad (16)$$

where A is the anchor point index set and the tolerance can be grouped together as $\epsilon = \{\epsilon_s, \epsilon_\theta, \epsilon_r, \epsilon_\phi\}$.

Then, the probability that a random representative ridge \mathbf{x}_a would match a random representative ridge \mathbf{x}_b is given by

$$\begin{aligned} p_\epsilon(\mathbf{x}) &= p(|\mathbf{x}_a - \mathbf{x}_b| \leq \epsilon | \Theta) \\ &= \int_{\mathbf{x}_a} \int_{|\mathbf{x}_a - \mathbf{x}_b| \leq \epsilon} p(\mathbf{x}_a | \Theta) p(\mathbf{x}_b | \Theta) d\mathbf{x}_a d\mathbf{x}_b \end{aligned} \quad (17)$$

where Θ is the set of parameters describing the distribution of the representative ridges.

The n PRC and specific n PRC with ridge information can be calculated by Eq.11, 13 and 14.

3.4 Experiments and Results

Parameters of the two fingerprint generative models introduced in Sections 2.2 and 2.3 were evaluated using the NIST fingerprint database. The NIST fingerprint database, NIST Special Database 4, contains 8-bit gray scale images of randomly selected fingerprints. Each print is 512×512 pixels with 32 rows of white space at the bottom of the print. The entire database contains fingerprints taken from 2000 different fingers with 2 impression of the same finger. Thus, there are a total of 2000 fingerprints using which the model has been developed. The number of components K for the mixture model was found after validation using k -means clustering.

Values of PRC p_ϵ are calculated using the formula introduced in Section 3. The tolerance is set at $\epsilon_s = \epsilon_r = 10$ pixels and $\epsilon_\theta = \epsilon_\phi = \pi/8$. For comparison,

Table 3. PRC for different fingerprint matches with varying m_1 (number of minutiae/ridges in fingerprint f_1), m_2 (number of minutiae/ridges in fingerprint f_2) and \hat{m} (number of matched minutiae/ridges)

| m_1 | m_2 | \hat{m} | Minutiae Only | | Minutiae and Ridge Points | |
|-------|-------|-----------|----------------------|----------------------|---------------------------|-----------------------|
| | | | p_ϵ | \hat{p}_ϵ | p_ϵ | \hat{p}_ϵ |
| 16 | 16 | 12 | 2.2×10^{-5} | 5.8×10^{-6} | 2.0×10^{-10} | 3.7×10^{-15} |
| 36 | 36 | 12 | 2.1×10^{-3} | 3.4×10^{-3} | 3.0×10^{-4} | 1.2×10^{-5} |
| 56 | 56 | 12 | 6.4×10^{-3} | 6.6×10^{-3} | 1.3×10^{-3} | 1.4×10^{-4} |
| 76 | 76 | 12 | 9.1×10^{-3} | 1.1×10^{-2} | 2.5×10^{-3} | 5.5×10^{-4} |

the empirical PRC $\hat{p}_\epsilon(\mathbf{x})$ was calculated with the same tolerance. To compute $\hat{p}_\epsilon(\mathbf{x})$, the empirical probabilities of matching a minutiae pair or ridge pair between imposter fingerprints are calculated first by

$$\hat{p}_\epsilon(\mathbf{x}) = \frac{1}{I} \sum_{i=1}^I \frac{\hat{m}_i}{m_i \times m'_i} \quad (18)$$

where I is the number of the imposter fingerprints pairs, \hat{m}_i is the number of matched minutiae or ridge pairs and m_i and m'_i are the numbers of minutiae or pairs in each of the two fingerprints. Then, the empirical PRC \hat{p}_ϵ can be calculated by Eq.10.

Both the theoretical and empirical PRCs are given in Table 3. The PRCs are calculated through varying numbers of minutiae or ridges in two randomly chosen fingerprint f_1 and f_2 and the number of matches between them. We can see that more minutiae or ridges the template and input fingerprint have, higher the PRC is.

Table 4. n PRCs with varying m and \hat{m} given $n = 100,000$ fingerprints

| No. of minutiae/ridges m | No of matches \hat{m} | Minutiae only | Minutiae and Ridge Points |
|----------------------------|-------------------------|--------------------------|---------------------------|
| | | $p(n)$ | $\hat{p}(n)$ |
| 46 | 46 | 5.0680×10^{-34} | 6.7329×10^{-49} |
| | 26 | 1.1842×10^{-3} | 1.7579×10^{-11} |
| 36 | 36 | 1.5551×10^{-22} | 7.5000×10^{-35} |
| | 16 | 1 | 3.4852×10^{-1} |
| 16 | 16 | 1.3464×10^{-1} | 4.460×10^{-8} |
| | 6 | 1 | 1 |

Based on the PRC value, n PRC can be computed. Table 4 shows the n PRCs in 100,000 fingerprints through varying number of minutiae or ridges in each fingerprint m and number of matches \hat{m} .

The specific n PRCs are also computed by (13) and given by Table 5. Here three fingerprints are chosen as query prints and they are shown in Figure 5. The first one is a full print in good quality, the second one is a full print in low quality and the third one is a partial print. The specific n PRCs are calculated through varying number of minutiae/ridges in each template fingerprint (m) and the number of matches (\hat{m}), assuming that the number of fingerprints in template database (n) is 100,000. The numbers of minutiae/ridges in 3 given query fingerprint m_f are 41, 26 and 13.



Fig. 5. Three specific fingerprints (from the same finger) used to calculate probabilities: (a) good quality full print F_1 , (b) low quality full print F_2 , and (c) partial print F_3 .

Table 5. Specific n PRCs for fingerprints in Fig. 5 with $n = 100,000$ fingerprints

| Fingerprint f | No. of m/r m_f | No. of m/r m | No. of matches \hat{m} | Minutiae only | Minutiae and Ridge Points |
|--------------------|---------------------|-------------------|-----------------------------|--------------------------|---------------------------|
| | | | | $p(n, f)$ | $p(n, f)$ |
| F_1 | 41 | 40 | 31 | 3.0327×10^{-78} | 8.5202×10^{-190} |
| | | 20 | 12 | 1.2995×10^{-18} | 2.6310×10^{-28} |
| F_2 | 26 | 20 | 12 | 1.1057×10^{-21} | 5.9308×10^{-45} |
| | | 10 | 4 | 8.3675×10^{-2} | 6.6823×10^{-4} |
| F_3 | 13 | 20 | 12 | 2.0588×10^{-24} | 1.4185×10^{-76} |
| | | 10 | 4 | 4.1707×10^{-2} | 1.2057×10^{-3} |

4 Summary

Generative models of individuality attempt to model the distribution of features and then use the models to determine the probability of random correspondence. We have proposed such models of fingerprint individuality for ridge flow,

minutiae and representative ridges. Individuality is evaluated in terms of three probability measures: probability of random correspondence (PRC) between two individuals, general probability of random correspondence (n PRC) between two individuals among a group of n individuals and specific probability of random correspondence (specific n PRC) which is the probability of matching a given individual among n individuals. We perform the individuality estimation on NIST4 dataset. As expected, we found that PRCs are reduced when more features are incorporated in the model. The proposed ridge information model offers a more reasonable and more accurate fingerprint representation. The results provide a much stronger argument for the individuality of fingerprints in forensics than previous generative models.

References

1. United States Court of Appeals for the Third Circuit: USA v. Byron Mitchell. No. 02-2859 (2003)
2. D. A. Stoney: Measurement of Fingerprint Individuality. Advances in Fingerprint Technology, H. Lee and R.E. Gaensslen, CRC Press (2001)
3. F. Galton: Finger Prints. McMillan, London (1892)
4. T. Roxburgh: Galton's work on the evidential value of fingerprints. Indian Journal of Statistics Vol.1, 62 (1933)
5. Henry, E.R.: Classification and Uses of FingerPrints. 54 Routledge & Sons, London (1900)
6. Trauring, M.: Automatic comparison of finger-ridge patterns. Nature, 197 (1963)
7. Pankanti, S. and Prabhakar, S. and A. K. Jain: On the individuality of fingerprints. IEEE Transactions on Pattern Analysis and Machine Intelligence, 24(8) (2002)
8. Dass, S.C. and Zhu, Y. and Jain, A. K.: Statistical Models for Assessing the Individuality of Fingerprints. Fourth IEEE Workshop on Automatic Identification Advanced Technologies, 3-9 (2005)
9. C. Su and S. N. Srihari: Generative Models for Fingerprint Individuality using Ridge Models. Proceedings of International Conference on Pattern Recognition, IEEE Computer Society Press (2008)
10. David R. Ashbaugh: Quantitative-Qualitative Friction Ridge Analysis: An Introduction to Basic and Advanced Ridgeology. CRC Press, Boca Raton, FL (1999)
11. C. Watson and M. Garris and E. Tabassi and C. Wilson and R. McCabe and S. Janet: User's Guide to NIST Fingerprint Image Software 2. 80-86, NIST (2004)
12. C. Bishop: Pattern Recognition and Machine Learning. Springer, New York (2006)
13. K. V. Mardia and P. E. Jupp: Directional Statistics. Wiley (2000)
14. Jiansheng Chen and Yiu-Sang Moon: A Minutiae-based Fingerprint Individuality Model. Computer Vision and Pattern Recognition, 2007. CVPR '07. (2007)
15. R.L. Vicent Hsu and Brian Martin: An Analysis of Minutiae Neighborhood Probabilities. Biometrics: Theory, Applications and Systems (2008)
16. Srihari, S. N. and S. Cha and H. Arora and S. J. Lee: Discriminability of Fingerprints of Twins. Journal of Forensic Identification Vol.58 1 109-127 (2008)
17. E. Tabassi and C. L. Wilson and C. I. Watson: Fingerprint Image Quality. NISTIR 7151, National Institute of Standards and Technology, August (2004)
18. Fang, G. and Srihari, S. N. and Srinivasan, H. and Phatak, P: Use of Ridge Points in Partial Fingerprint Matching. Biometric Technology for Human Identification IV, 65390D1-65390D9 SPIE (2007)